

Therapies in Metastatic Non-small-cell Carcinoma

a report by

Prina R Ruparelia and Stephen G Spiro

Department of Thoracic Medicine, University College Hospital, London

DOI: 10.17925/EOH.2007.0.1.46

Lung cancer is the most lethal cancer in men and women in the western world. Eighty per cent of cases are non-small-cell lung carcinoma (NSCLC) whose subtypes include squamous cell carcinoma, adenocarcinoma and large-cell carcinoma. Of these, approximately 80% will present with either locally advanced or metastatic disease – i.e. stage IIIB or IV disease – and are unlikely to be cured.

Treatment options for these patients include best supportive care (BSC), chemotherapy or a combination of radiotherapy and chemotherapy. NSCLC are less chemosensitive than small-cell carcinoma and complete responses are rare, but nevertheless chemotherapy provides some improvements in quality of life (QoL) and life expectancy compared with BSC, particularly in better performance status (PS) patients. This article will discuss the medical treatment of NSCLC, with some attention to the newer epidermal growth factor (EGFR) antagonists.

After chemotherapy, overall one-year survival in advanced disease was 30% compared with best supportive care, where one-year survival was only 17%.

Chemotherapy versus Best Supportive Care

Chemotherapy became a serious entity for NSCLC only in the mid-1990s, particularly after a meta-analysis by the Non-small Cell Lung Cancer Collaborative Group who looked at all randomised controlled trials (RCTs) of chemotherapy or BSC in all treatment situations (i.e. surgery, radiotherapy or BSC).¹ In trials of chemotherapy versus BSC in advanced disease, nine RCTs showed that chemotherapy conferred a small survival advantage of six to eight weeks. However, in these retrospective analyses there were no QoL data. Chemotherapy regimes may have considerable side effects, thus it is important to take this into account. A balance will inevitably have to be struck between QoL and the effects of chemotherapy, which may be influenced by any additional morbidity of these regimes. After chemotherapy, overall one-year survival in advanced disease was 30% compared with BSC, where one-year survival was only 17%.

Chemotherapy Regimens

The NSCLC Collaborative Group meta-analysis showed that only cisplatin-containing regimens were effective, but the advantages were small.¹ Although a number of different regimens of these second-generation compounds have been used, there is little to choose between them.

However, several drugs in combination were better than single-agent therapy, with better response rates and prolonged survival, at the cost of increased toxicity with combination therapy. Common second-generation chemotherapy regimens include mitomycin, ifosfamide and cisplatin (MIC) and mitomycin, vinblastine and cisplatin (MVP). The effect of MIC versus BSC was studied by Cullen (1999) in 351 patients, demonstrating a prolongation in median survival of seven weeks.² Smith et al. studied the effect of another common regime, and in particular the duration of chemotherapy, by administering either three or six cycles of MVP.³ The authors concluded that there was no benefit in continuing chemotherapy beyond three cycles in terms of median survival, one-year survival and QoL. These regimes were used in over 70% of the patients in the Big Lung trial, which compared BSC with four different chemotherapy regimes and importantly demonstrated a survival benefit with no detrimental effect in terms of QoL for the addition of chemotherapy.⁴ This advantage had been hinted at, for example, by Cullen et al., who compared four courses of MIC with BSC and demonstrated an increase in median survival of two months.² The overall response rate to chemotherapy was 31%, with a complete response of 2% and a partial response of 29%. Only a small subset of patients entered a QoL study and this showed an improvement in the symptom scores as measured by the European Organisation for Research and Treatment of Cancer (EORTC QLC-LC13). However, the largest study of this type was the Big Lung trial, with nearly 400 patients recording QoL.

Third-generation agents are more expensive, but have the advantage of better tolerance and can be given as outpatient treatment.

Over the last decade, new drugs have emerged, including gemcitabine, vinorelbine and the taxols. These third-generation agents are more expensive, but have the advantage of better tolerance and can be given as outpatient treatment. However, once again, there was little to choose between regimens such as carboplatin/gemcitabine, carboplatin/vinorelbine or carboplatin/paclitaxel, with a median survival of about nine to 10 months and an increase to 35–40% of patients alive at one year (Schiller, 2002).⁵ Docetaxel has also emerged as an effective single agent, with 15–20% responding when used as second-line treatment at relapse.⁶

In the UK, carboplatin/gemcitabine has become the favoured first-line

treatment after the study of Rudd et al., which compared gemcitabine and carboplatin (GC) with the older regimen of MIC.⁷ GC conferred a median survival of 10 months compared with a median survival of 7.6 months for MIC with one-year survivals of 40% and 30%, respectively. The response rates were similar for the two regimens: 42% for GC and 41% for MIC. GC had fewer side effects and resulted in fewer hospital admissions, and there was a highly significant advantage for most QoL variables studies for the newer regimen.

The mean age at presentation with lung cancer in the UK is 68 years of age, with 40% of patients presenting at 70 years or above.

Chemotherapy in the Elderly Patient

The mean age at presentation with lung cancer in the UK is 68 years of age, with 40% of patients presenting at 70 years or above. Most of the phase III RCTs tend to recruit younger patients – in most trials, just 10% of patients are over 70 years of age. Hence, prospective data on tolerance and responses in the elderly tend to be retrospective data from large trials, which did not specifically set out to determine how the elderly would fare with chemotherapy. It is clear that more elderly patients either refuse or are not offered chemotherapy.⁸ Earle et al. showed that, in 2000, only 22% of patients older than 65 years received chemotherapy, and the Surveillance, Epidemiology and End Results data from the US (collected between 1988 and 1995) showed that only 32% of patients over 65 years of age received chemotherapy.⁹

There have been some specific trials of chemotherapy in elderly patients with NSCLC. Langer et al. studied the effects of three different regimens: cisplatin/etoposide, high-dose paclitaxel/cisplatin and low-dose paclitaxel/cisplatin.¹⁰ The two taxol arms had a better response, with the median survival being 10 and 9.5 months in the taxol arms versus 7.6 months in the non-taxol arm. QoL changes were similar to studies in younger patients. Although the response rate, toxicity and survival were similar to younger patients, there was more co-morbidity, leukopaenia and neuropsychiatric toxicity in these older patients. The MILES study looked at the effect of combination chemotherapy versus a single agent, vinorelbine.¹¹ Six hundred and ninety-eight patients were entered and demonstrated that combination chemotherapy did not provide any survival benefit compared with vinorelbine alone.¹² Frasci et al., however, showed that combination therapy produced a superior median survival, but the chemotherapy doses used were different in the two studies, making comparisons difficult. The Elderly Lung Cancer Vinorelbine Italian Study compared vinorelbine as a single agent with BSC.¹³ The study was terminated early due to poor recruitment. However, in the 191 patients recruited, a prolongation in median survival of seven weeks was demonstrated with chemotherapy. Patients who received vinorelbine performed better on functional assessments but scored worse on toxicity. These studies were performed on patients with a PS of one or two; the majority of patients had a PS of one. In general, good PS patients tolerate and respond better to the chemotherapy; some PS 2 patients can also do well, but in general they

suffer more toxicity than those with PS 0–1, and the progression-free interval is shorter.

Trials of chemotherapy in elderly patients have provided conflicting results, with better median survivals than with BSC, and with a suggestion that single-agent chemotherapy may be less toxic than combination therapy and confers a small survival benefit.¹⁴

Chemotherapy and Radiotherapy

Local radiotherapy is a useful tool in the palliation of lung cancer. Radical radiotherapy is used in the treatment of stage IIIA and IIIB disease, and most studies report a median survival of eight to 12 months. There has been a lot of interest as to whether a combination of radiotherapy and chemotherapy provides an additional survival benefit in locally advanced NSCLC. Many trials have looked at this question.¹ A meta-analysis by the NSCLC collaborative group looked at 22 RCTs including a total of 3,033 patients randomised to have chemotherapy alone or combination chemotherapy and radiotherapy. The patients who received combination treatment demonstrated a prolongation in their median survival of seven to eight weeks.¹⁵ Furuse et al. have taken this question further to determine whether concurrent chemoradiotherapy or sequential chemotherapy and radiotherapy is more advantageous. Patients were randomised to receive either concurrent therapy with mitomycin, vindesine, cisplatin and radiotherapy or sequential chemotherapy and radiotherapy. Concurrent chemoradiotherapy significantly enhanced median survival (16.5 versus 13.3 months). Trials are ongoing concerning the timing of chemotherapy and radiotherapy in NSCLC.

Epidermal growth factor inhibitors have been demonstrated on the cell surface of 60–90% of non-small-cell lung cancers.

Non-small-cell Lung Carcinoma and Performance Status

PS is an independent prognostic factor. Patients with a PS of two demonstrate lower response rates to chemotherapy and shorter time to treatment failure, have higher toxicity and tolerate fewer courses of chemotherapy. Many trials have looked at the effects of chemotherapy versus best supportive care. The trials all used different chemotherapy regimens; however, the consensus opinion at present is to use single-agent therapy with gemcitabine, vinorelbine or a taxane. The alternative regime is carboplatin-based combination chemotherapy.

Epidermal Growth Factor Inhibitors

EGFR have been demonstrated on the cell surface of 60–90% of non-small-cell lung cancers. Downstream pathways stimulated by EGFR phosphorylation result in cancer cell proliferation and inhibition of apoptosis, among other effects. Drugs have been developed that block this receptor. Gefitinib was the first agent developed. The mechanism of action of gefitinib is to block EGFR tyrosine kinases, reducing cell proliferation.^{16,17} The Iressa Dose Evaluation in Advanced Lung Cancer (IDEAL 1 and 2) trials gave gefitinib at 250 and 500mg; there was no

placebo arm in this study. Gefitinib was given to patients who had been previously treated with two or more chemotherapy regimes and had relapsed. There were approximately 200 patients in each trial. Symptom improvement occurred in both treatment groups and differences between the two doses were not significantly different. The radiographical response rate was also similar between the two dosing groups, with a partial response rate of 13 and 9% in the 250 and 500mg groups, respectively, although it was generally lower in the more heavily pre-treated patients. These uncontrolled studies provided evidence supporting the launch of gefitinib for general use in pre-treated patients at relapse.

However, a larger trial comparing gefitinib at relapse to placebo in more than 1,600 patients (the Iressa Survival Evaluation in Lung cancer (ISEL) trial) showed no survival benefit for the addition of gefitinib.¹⁸ However, subgroup analyses showed a survival benefit in Asian patients, females, non-smokers and the adenocarcinoma cell type. Another mode of timing was studied in the Iressa NSCLC Trial Assessing Combination Treatment

another EGFR inhibitor, erlotinib.²⁰ It was used in patients who had had one or two previous chemotherapy regimes, and the trial included a placebo arm. This study demonstrated a prolonged median survival of two months with erlotinib compared with placebo and 31% of patients were alive at one year in comparison with 22% in the placebo arm.

Developments in targeted therapy may be increasingly important in the future...

Third-generation combination chemotherapy achieves a 30–40% partial response rate...

(INTACT) trials, where gefitinib was randomised to be given at the beginning of treatment with a platinum-based regimen of either cisplatin and gemcitabine or carboplatin and paclitaxel.¹⁹ These two trials demonstrated no benefit in terms of median survival, time to progression, response rate or overall survival by the addition of these drugs to standard chemotherapy regimes.

Shepherd et al. performed a large RCT in relapsed advanced NSCLC with

Interestingly, multivariate analyses again demonstrated that Asian origin, female gender and non-smoking were independent variables for better survival with erlotinib. Analysis of the histological subtype identified that patients with adenocarcinoma were more likely to experience a benefit. Thus, current evidence shows that the use of these agents in combination with first-line chemotherapy is not beneficial, but that, in relapsed disease, these agents provide a median survival advantage of up to eight weeks. Research is now attempting to predict which patients are most likely to respond to these agents and why, so that they may become truly targeted therapy – a claim that seems premature at present.

Conclusion

Several chemotherapy agents demonstrate activity in NSCLC. There are several regimes available, although no regimen stands out. Third-generation combination chemotherapy achieves a 30–40% partial response rate and prolongs median survival by three to four months compared with BSC, with about 35% of patients alive at one year. Modern chemotherapy has fewer toxic effects compared with the older regimens of the early 1990s. Developments in targeted therapy may be increasingly important in the future, as chemotherapy seems to have reached a plateau. ■

1. Non-small cell lung cancer collaborative group, Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials, *BMJ*, 1995;311(7010):899–909.
2. Cullen MH, Billingham LJ, Woodroffe CM, et al., Mitomycin, ifosfamide and cisplatin in unresectable non small cell carcinoma: effects on survival and quality of life, *J Clin Oncol*, 1999;17(10):3188–94.
3. Smith IE, O'Brien ME, Talbot DC, et al., Duration of chemotherapy in advanced non-small cell cancer: a randomised trial of three versus six courses of mitomycin, vinblastine and cisplatin, *J Clin Oncol*, 2001;19(5):1336–43.
4. Rudd RM, Gower NH, Spiro SG, et al., Gemcitabine plus carboplatin versus mitomycin, ifosfamide, and cisplatin in patients with stage IIIB or IV non-small cell lung cancer: a phase III randomised study of the London lung cancer group, *J Clin Oncol*, 2005;23(1):142–53.
5. Brown J, Thorpe H, Napp V, et al., Assessment of quality of life in the supportive care setting of the big lung trial in non-small cell lung carcinoma, *J Clin Oncol*, 2005;23(30):7417–27.
6. Schiller J, Harrington D, Belani CP, et al., Comparison of four chemotherapy regimens for advanced non-small lung cancer, *N Eng J Med*, 2002;346(2):92–8.
7. Roszkowski K, Pluzanska A, Krzakowski M, et al., A multicenter randomised phase III study of docetaxol plus best supportive care versus best supportive care in chemotherapy naïve with metastatic or non-resectable localised non-small cell lung cancer (NSCLC), *Lung Cancer*, 2000;27 (3):145–57.
8. Earle CC, Vendetti LN, Neumann PJ, et al., Who gets chemotherapy for metastatic lung cancer?, *Chest*, 2000;117(5):1239–46.
9. SEER Surveillance and epidemiology, <http://seer.cancer.gov>.
10. Langer CJ, Manola J, Bernardo P, Cisplatin-based therapy for elderly patients with advanced non-small cell cancer: implications of Eastern Cooperative Oncology group 5922 a randomised trial, *J Natl Cancer Inst*, 2002;94(3):173–81.
11. Gridelli C, Perrone F, Gallo C, et al., Chemotherapy for elderly patients with advanced non-small-cell lung cancer: the Multicenter Italian Lung Cancer in the Elderly Study (MILES) phase III randomized trial, *J Natl Cancer Inst*, 2003;95 (5):62–72.
12. Frasci G, Lorusso V, Panza N, et al., Gemcitabine plus vinorelbine versus vinorelbine alone in elderly patients with advanced non-small cell lung cancer, *J Clin Oncol*, 2000;18(13): 2529–36.
13. Gridelli C, The ELVIS trial: a phase III study of single agent vinorelbine as first line treatment in elderly patients with advanced non-small cell lung cancer, Elderly Lung Cancer Vinorelbine Italian Study, *Oncologist*, 2001;6(Suppl. 1):4–7.
14. Lilenbaum RC, Herndon JE II, List MA, et al., Single-agent versus combination chemotherapy in advanced non-small-cell lung cancer: the cancer and leukemia group B (study 9730), *J Clin Oncol*, 2005;23(1):190–96.
15. Furuse K, Fukuoka M, Kawahara M, et al., Phase III study of concurrent versus sequential radiotherapy in combination with mitomycin, vindesine and cisplatin in unresectable stage III non-small-cell lung cancer, *J Clin Oncol*, 1995;17:2692–9.
16. Fukuoka M, Yano S, Giaccone G, et al., Multi-institutional randomised phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer (the IDEAL trial), *J Clin Oncol*, 2003;21:2237–46.
17. Kris MG, Natale RB, Herbst RS, et al., Efficacy of gefitinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase in symptomatic patients with non-small-cell lung cancer, *JAMA*, 2003;290:2149–58.
18. Thatcher N, Chang A, Parikh P, et al., Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell lung cancer: results from a randomised, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer), *Lancet*, 2005;366(9496): 1527–37.
19. Giaccone G, Herbst RS, Manegold C, et al., Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial-INTACT 1, *J Clin Oncol*, 2004;22:785–94.
20. Shepherd FA, Pereira JR, Ciuleanu T, et al., Erlotinib in previously treated non-small-cell lung cancer, *New Eng J Med*, 2005;353:123–32.