

# The Harmful Effects of Iron Overload in Patients with Myelodysplastic Syndrome

a report by

Andrea Toma<sup>1</sup> and François Dreyfus<sup>2</sup>

1. Clinical Haematology Department, Pitié Salpêtrière Hospital, Paris;

2. Professor of Medicine, University of Paris Descartes–Paris V, and Head, Haematology Unit, Hospital Cochin, Paris

DOI: 10.17925/EOH.2008.02.1.75

Myelodysplastic syndrome (MDS) refers to a group of clonal haemopoietic disorders characterised by ineffective haemopoiesis, bone marrow dysplasia and an increased risk of transformation to acute myelogenous leukaemia (AML). A number of prognostic scoring systems have been developed to predict the outcomes of MDS. The International Prognostic Scoring System (IPSS), which uses karyotype, percentage of bone marrow blasts and the number of cytopenia, is the most widely used system to predict survival and the risk of progression to AML.<sup>1</sup> Low- and high-risk MDS is characterised by low, int-1, int-2 and a high IPSS risk.

The majority of patients with MDS are anaemic. Up to 90% of patients will require regular transfusions. In 50–60% of patients, anaemia is severe (haemoglobin level below 10g/dl)<sup>1</sup> and is associated with a decreased physical performance and poor quality of life,<sup>2</sup> along with increased cardiac morbidity and mortality.<sup>3–5</sup>

The aim of blood transfusion therapy is to maintain quality of life and prevent anaemia-related morbidity and mortality. Several studies have now shown that transfusion dependency has a negative impact on the survival of patients with MDS.<sup>3,6</sup> A number of factors may contribute to the decreased survival of transfusion-dependent patients. It has been clearly shown that mortality correlates with higher serum ferritin levels in patients with MDS, probably as a consequence of iron overload.<sup>3,7</sup>

## Mechanisms of Iron Overload and Toxicity in Myelodysplastic Syndrome Patients

Several mechanisms play a role in iron loading in patients with MDS, who start accumulating iron even before they are in need of regular blood transfusions. At this stage of the disease, iron accumulation is probably the result of ineffective erythropoiesis, which stimulates increased iron absorption by the small intestine. The mechanism of iron accumulation at this stage is unclear, but downregulation of hepcidin, which is an important modulator of serum ferritin levels, is probably important.<sup>8,9</sup> In the GFM registry, before transfusions begin, the serum ferritin level has usually risen to about 500–600µg/l and seldom exceeds this value.

The main cause of iron overload is transfusion therapy. Each unit of blood transfused delivers 200–250mg of iron. After only 20 units of transfused blood, 4,000–5,000mg of iron will have been delivered to the body. At this iron-loading, the serum ferritin level rises to about 1,000µg/l,<sup>3</sup> and the capacity of transferrin to bind iron is exceeded. Reactive non-transferrin-bound iron (NTBI) is then generated,<sup>10</sup> favouring oxidative DNA damage and apoptosis.<sup>11,12</sup> NTBI levels are significantly elevated compared with normal levels even in non-transfused patients with MDS. Those patients with higher NTBI levels show a higher level of apoptosis in the bone marrow.<sup>13</sup>

## Consequences of Iron Overload in Myelodysplastic Syndrome Patients

In 1981 it was reported that patients with refractory or aplastic anaemia, who had received an average of 120 blood transfusions, had seven to 26 times the normal amount of iron in the liver and typically showed focal portal fibrosis.<sup>14</sup> Although left ventricular cardiac function was impaired in only the most heavily transfused patients or in those with co-existing coronary artery disease<sup>15,16</sup> (see *Table 1*), all patients had glucose intolerance associated with a significantly reduced insulin output when compared with controls. This pattern of organ involvement resembles that encountered in idiopathic haemochromatosis.<sup>14</sup>

Two other retrospective studies have highlighted the contribution of iron overload to the morbidity and mortality associated with MDS. More than 20 of 46 patients with MDS who had received more than 50 units of blood showed signs of heart failure, sometimes with cardiac arrhythmias. Fourteen patients died of congestive heart failure secondary to haemochromatosis in less than four years.<sup>17</sup> In another smaller study, 34 of 43 patients with the 5q-subtype of MDS were transfusion-dependent at diagnosis, and 40 patients were eventually transfusion-dependent.<sup>18</sup> Haemosiderosis was diagnosed by iron studies in 12 patients and was associated with congestive heart failure or exertion intolerance, abnormal liver function tests, cirrhosis, diabetes and skin changes.



Andrea Toma is a Doctor in the Clinical Haematology Department of Pitié Salpêtrière Hospital in Paris. Her major research interests in myelodysplastic syndrome (MDS) include iron metabolism and the immunological abnormalities observed in MDS. She is a member of the Groupe Français des Myélodysplasies (GFM). Dr Toma studied medicine at the University of Paris VI and received her MD from the University of Paris Descartes – Paris V. She began specialist training in haematology in 2006. She received her licence of biological and medical science in immunology in 1999, and completed her Masters in 2000 at the Pasteur Institute and a research fellowship in immunology in 2003.



François Dreyfus is a Professor of Medicine at the University of Paris Descartes–Paris V and Head of the Haematology Unit at Hospital Cochin in Paris, which is recognised as a centre of excellence by the Myelodysplastic Syndromes Foundation. His major research interests include epidemiology and pathogenesis, particularly the mechanisms of apoptosis in myelodysplastic syndrome. His work, particularly on myelodysplastic syndrome, is recognised internationally. Professor Dreyfus is a member of the

Société Française Hématologie and Myelodysplastic Syndromes Foundation and Vice President of the Groupe Francophone des Myélodysplasies (GFM). Professor Dreyfus is the author or co-author of more than 160 papers published in peer-reviewed journals including *Blood*, the *British Journal of Haematology*, *Haematologica*, the *Journal of Clinical Oncology* and *Leukaemia Research*. He studied medicine at the University of Diderot–Paris VII and received his MD from the University of Paris Descartes–Paris V. He began specialist training in haematology in 1992.

E: francois.dreyfus@cch.ap-hop-paris.fr

**Table 1: Incidence of Cardiac Events over a Three-year Period Among 705 patients with Myelodysplastic Syndrome in the US Medicare Population**

	Number of Patients (%)
Any cardiac-related event	522 (74)
Myocardial infarction	142 (29)
Congestive heart failure	344 (48)
Arrhythmias	374 (53)
Other cardiac events	415 (58)

Data from Goldberg et al., 2007.<sup>19</sup>

**Table 2: Eligibility Criteria for Iron Chelation Therapy in Patients with Myelodysplastic Syndrome, According to the Nagasaki Consensus Statement<sup>32</sup>**

Transfusion Dependency
Serum ferritin level >1,000–2,000µg/litre or other evidence of significant tissue iron overload
Patients with low-risk MDS
*IPSS low or intermediate-1 category
*WHO-type refractory anaemia (RA), RA with ringed sideroblasts (RARS) and 5q- syndrome
Documented stable disease
No co-morbidities that severely limit prognosis
Candidates for allografting

Data from 374 patients with MDS show that once patients with MDS develop a need for regular transfusions, their probability of survival decreases (hazard ratio for death 1/58 with a probability of 0.005).<sup>6</sup> In a survey of 467 patients with MDS followed from diagnosis to death or progression to leukaemia, 51% of non-leukaemic deaths were due to cardiac failure and 8% were due to hepatic cirrhosis. Other related complications of MDS leading to death include haemorrhage and infection.<sup>3</sup> Cardiac failure was a significantly more common cause of death among MDS patients who were transfusion-dependent than among those not receiving regular transfusions. Iron overload was a major contributor to this increased mortality.

In a retrospective Japanese study of 292 patients with MDS and other transfusion-dependent anaemias, cardiac and hepatic dysfunction were present in 21.9 and 84.6% of patients studied, respectively.<sup>19</sup> Among the patients who died, 24.0% had cardiac failure, 6.7% had hepatic failure and 97% of those who died had serum ferritin levels above 1,000µg/l.

In a large retrospective study of 840 consecutive patients with MDS, heart failure (28 versus 18%;  $p=0.001$ ) and cardiac death (69 versus 55%;  $p=0.03$ ) were significantly more common in transfusion-dependent than in non-transfusion-dependent patients.<sup>20</sup> In a Cox analysis with time-dependent co-variables, the risk of non-leukaemic death (hazard ratio 2/12;  $p<0.001$ ), heart failure (hazard ratio 1/34;  $p=0.03$ ) and cardiac death (hazard ratio 2/99;  $p=0.01$ ) were all increased in transfusion-dependent patients. The development of secondary iron overload significantly increased the risk of non-leukaemic death (hazard ratio 1/15;  $p<0.001$ ) and specifically increased the risk of developing heart failure (hazard ratio 1/17;  $p<0.001$ ).

Transfusion dependency may be a reliable indicator of the evolution of the disease. It must be recognised that the limited survival of patients with MDS may be, at least partly, caused by a deterioration of their stem cell disorder (for example the progression of MDS to AML) and the presence of co-morbidities.<sup>21</sup> This is in addition to the direct effect of iron overload as a result of blood transfusions.

Myocardial accumulation of iron has not yet been extensively studied in patients with MDS. The results that have been obtained are contradictory. Recent myocardial T2\* magnetic resonance imaging (MRI) studies suggested that iron-loading of the heart can occur after 75–100 units of blood have been transfused.<sup>22</sup> The value of T2\* is only modified by iron overload independent of coronary disease. Two recent studies using T2\* MRI failed to show an iron accumulation in the hearts of patients with MDS after 63–90 units of blood. Iron accumulation in the liver was shown in all of the patients studied.<sup>23,24</sup> A recent report from an Italian group study confirms the relationship between cardiac overload and transfusion burden (of more than 100 blood units).<sup>25</sup> It was a short and heterogeneous series (some patients received chelation therapy, others did not), which may have contributed to the findings. Increased NTBI in patients with MDS may cause direct myocardial tissue damage. This cannot always be shown by MRI, which visualises iron storage.<sup>26</sup> It has been suggested that there may be a long latency period between iron accumulation in the liver and iron overload in the hearts of patients with MDS.<sup>15</sup> This is supported by a more recent T2\* MRI series.<sup>27</sup>

## Outcomes of Chelation Therapy in Myelodysplastic Syndrome

The positive impact of iron chelation therapy on survival has been clearly shown in patients with  $\beta$ -thalassaemia major, in whom maintenance of serum ferritin levels below 2,500µg/l was the major determinant of cardiac disease-free survival.<sup>28</sup> Although there are few studies of the efficacy of iron chelation therapy in patients with MDS, evidence is slowly accumulating that efficient iron chelation could have a positive impact on survival in patients with MDS.

A retrospective study on 178 patients with MDS indicated that iron chelation therapy was significantly associated with improved survival in patients with low- or intermediate-1-risk MDS.<sup>29</sup> The median overall survival rate was not reached at 160 months in patients who received iron chelation therapy. It was only 40 months in those who did not receive iron chelation. Significantly more patients who received iron chelation therapy survived up to four years (80 versus 44%;  $p<0.03$ ). Serum ferritin levels fell significantly in patients who received iron chelation therapy, and increased significantly in those who did not. In another retrospective study in 292 patients with MDS or other transfusion-dependent anaemias, effective iron chelation therapy with deferoxamine (daily or continuous chelation therapy) resulted in improved levels of serum ferritin, liver enzymes and fasting blood sugar levels.<sup>19</sup> Intermittent deferoxamine therapy conferred no benefit.

These retrospective findings need to be confirmed by prospective studies. In a prospective cohort study that enrolled 170 patients (five of whom were lost to follow-up) with MDS receiving regular blood transfusions, iron chelation therapy significantly improved the median overall survival further to diagnosis from 51 to 115 months ( $p<0.0001$ ).<sup>30</sup> After adjustment for other prognostic factors, such as sex, age, IPSS category and transfusion requirement, the survival difference remained significant. The main prognostic factor of survival in Cox analysis is correct iron chelation. Most importantly the difference of survival is due to non-leukaemic death.

## Eligibility for Iron Chelation Therapy

Current guidelines for the management of iron chelation therapy in patients with MDS are based on non-randomised trials, expert opinion and the experience of iron chelation in patients with  $\beta$ -thalassaemia

major.<sup>31–35</sup> These guidelines converge towards a consensus on the subgroups of patients with MDS who should receive chelation therapy. The Italian guidelines recommend that adult patients with MDS who have previously received more than 50 units of blood and with an expected lifespan longer than six months should receive iron chelation therapy.<sup>31</sup> The UK guidelines recommend that iron chelation should be considered once a patient has received 5g of iron or approximately 25 units of blood, but only if long-term transfusional therapy is likely.<sup>32</sup> The most recent National Comprehensive Cancer Network (NCCN) practice guidelines strongly recommend that iron chelation therapy be considered in patients who have received 20–30 units of blood and for whom ongoing blood transfusions are anticipated.<sup>35</sup>

The Nagasaki consensus recommends starting iron chelation therapy when serum ferritin levels reach 1,000–2,000µg/l, dependent on the transfusion rate.<sup>32</sup> Expert opinion is now of the view that a serum ferritin level of 1,000µg/l is a suitable threshold for starting iron chelation therapy. Chelation therapy should continue for as long as transfusion therapy continues and as long as iron overload remains clinically relevant.<sup>31–35</sup> The aim is to reach and maintain a serum ferritin level below 1000µg/l.

The MDS Foundation guidelines recommend that iron chelation therapy should be initiated when the serum ferritin reaches 1,000µg/l, and/or after the patient has received two units of RBCs per month for at least one year. Iron chelation therapy should be proposed to patients with an anticipated survival superior to one year, without co-morbidities, limiting prognosis or candidates for allograft.<sup>34</sup>

The guidelines all agree that RBC transfusions are clinically beneficial to treat symptomatic anaemia in MDS. Patients with low-risk MDS receiving transfusions are the most likely to benefit from iron chelation therapy. The NCCN guidelines emphasise that iron chelation should, in particular, be considered for patients with low-risk MDS whose clinical course suggests the need for continuing red blood cell transfusions along with those with concurrent cardiac or hepatic dysfunction.<sup>35</sup> These guidelines recognise that patients with MDS may have additional factors that make them more vulnerable to the toxic effects of iron overload.

The Nagasaki consensus statement provides more detailed eligibility criteria. Patients with refractory cytopenia with multilineage dysplasia (RCMD) or refractory sideroblastic cytopenia with multilineage dysplasia (RSCMD), coupled with a low IPSS score, are also good candidates for iron chelation therapy. These patients are likely to survive for about five years. Patients with RCMD, RSCMD and an IPSS intermediate-1 score with a life expectancy of about three years may also benefit, though the indication is less clear.<sup>36</sup>

The MDS patient tends to have multiple clinical problems. The causes of decreased survival are likely to be multifactorial and not always related to transfusions. Retrospective and prospective evidence is accumulating on the deleterious role of iron in MDS especially in liver and heart dysfunctions. More prospective studies are difficult to interpret. Current guidelines for iron chelation therapy in patients with MDS who have a reasonable survival prognosis are largely based on clinical experience. ■

- Greenberg P, Cox C, LeBeau MM, et al., International scoring system for evaluating prognosis in myelodysplastic syndromes, *Blood*, 1997;89:2079–88.
- Penninx BW, Guralnik JM, Onder G, et al., Anemia and decline in physical performance among older persons, *Am J Med*, 2003;115:104–10.
- Malcovati L, Porta MG, Pascutto C, et al., Prognostic factors and life expectancy in myelodysplastic syndromes classified according to WHO criteria: a basis for clinical decision making, *J Clin Oncol*, 2005;23:7594–7603.
- Wu WC, Rathore SS, Wang Y, et al., Blood transfusion in elderly patients with acute myocardial infarction, *N Engl J Med*, 2001;345:1230–36.
- Oliva EN, Dimitrov BD, Benedetto F, et al., Hemoglobin level threshold for cardiac remodeling and quality of life in myelodysplastic syndrome, *Leuk Res*, 2005;29:1217–19.
- Cazzola M, Malcovati L, Myelodysplastic syndromes-coping with ineffective hematopoiesis, *N Engl J Med*, 2005;352:536–8.
- Malcovati L, Della Porta MG, Cazzola M, Predicting survival and leukaemic evolution in patients with myelodysplastic syndrome, *Haematologica*, 2006;91:1588–90.
- Wood JC, Enriquez C, Ghugre N, et al., Physiology and pathophysiology of iron cardiomyopathy in thalassemia, *Ann N Y Acad Sci*, 2005;1054:386–95.
- Winder A, Lefkowitz R, Ghoti H, et al., Urinary hepcidin excretion in patients with myelodysplastic syndrome (MDS) and myelofibrosis (MF), *Blood*, 2006;108: abstract 2620.
- Brittenham GM, Olivieri NF, Roualt TA, Iron physiology and iron overload, American Society of Hematology Education Programme, 1996:177.
- Grootveld M, Bell JD, Halliwell B, et al., Non transferrin-bound iron in plasma or serum from patients with idiopathic hemochromatosis, *J Biol Chem*, 1989;264:4417.
- Breuer W, Greenberg E, Cabantchik ZI, Newly delivered transferrin iron and oxidative cell injury, *FEBS Lett*, 1997;403:213.
- Corteletti A, Cattaneo C, Cristiani S, et al., Non-transferrin-bound iron in myelodysplastic syndromes: a marker of ineffective erythropoiesis?, *Hematol J*, 2000;1:153–8.
- Schafer AI, Cheron RG, Dluhy R, et al., Clinical consequences of acquired transfusional iron overload in adults, *N Engl J Med*, 1981;304:319–24.
- Buja LM, Roberts WC, Iron in the heart. Etiology and clinical significance, *Am J Med*, 1971;51:209–21.
- Goldberg SL, Moody-Patel N, Warnock N, Incidence of myelodysplastic syndromes in the United States Medicare population, *Leuk Res*, 2007;31(Suppl. 1):S103.
- Jaeger M, Aul C, Söhngen D, et al., Sekundäre Hämochromatose bei polytransfundierten Patienten mit Myelodysplastischen Syndromen, *Beits Infusionssther*, 1992;30:464–8.
- Mathew P, Tefferi A, Dewald GW, et al., The 5q- syndrome: a single-institution study of 43 consecutive patients, *Blood*, 1993;81:1040–45.
- Takatoku M, Uchiyama T, Okamoto S, et al., Retrospective nationwide survey of Japanese patients with transfusion-dependent MDS and aplastic anaemia highlights the negative impact of iron overload on morbidity/mortality, *Eur J Haematol*, 2007;78:487–94.
- Della Porta MG, Malcovati L, Travagino E, et al., A prognostic model for predicting the impact of comorbidities on survival of patients with myelodysplastic syndromes, *Blood*, 2007;(110): abstract 2453.
- Deschler B, de Witte T, Mertelsham R, Lübbert M, Treatment decision-making for older patients with high-risk myelodysplastic syndrome or acute myeloid leukaemia: problems and approaches, *Haematologica*, 2006;91:1513–22.
- Jensen PD, Jensen FT, Christensen T, et al., Evaluation of myocardial iron by magnetic resonance imaging during iron chelation therapy with deferoxamine: indication of close relation between myocardial iron content and chelatable iron pool, *Blood*, 2003;101:4632–9.
- Chacko J, Pennell DJ, Tanner MA, et al., Myocardial iron loading by magnetic resonance imaging T2\* in good prognostic myelodysplastic syndrome patients on long-term blood transfusions, *Br J Haematol*, 2007;138:587–93.
- Konen E, Ghoti H, Goitein O, et al., No evidence for myocardial iron overload in multitransfused patients with myelodysplastic syndrome using cardiac magnetic resonance T2\* technique, *Am J Hematol*, 2007;82:1013–16.
- di Tucci AA, Deplano GMS, et al., Myocardial iron overload assessment by T2\* magnetic resonance imaging in adult transfusion dependent patients with acquired anemias, *Haematologica*, 2008;93:1385–8.
- Anderson LJ, Holden S, Davis B, et al., Cardiovascular T2-star (T2\*) magnetic resonance for the early diagnosis of myocardial iron overload, *Eur Heart J*, 2001;22:2171–9.
- Glanville J, Eleftheriou P, Porter J, MRI evidence of cardiac iron accumulation in myelodysplasia and unusual anaemias, *Blood*, 2006;321:1255–8.
- Olivieri NF, Nathan DG, MacMillan JH, et al., Survival in medically treated patients with homozygous beta-thalassemia, *N Engl J Med*, 1994;331:574–8.
- Leitch HA, Improving clinical outcome in patients with myelodysplastic syndrome and iron overload using iron chelation therapy, *Leuk Res*, 2007;31(Suppl. 3):S7–S9.
- Rose C, Brechignac S, Vassilief D, et al., Positive impact of iron chelation therapy (CT) on survival in regularly transfused MDS patients. A prospective analysis by the GFM, *Blood*, 2007;110: abstract 249.
- Alessandrino EP, Amadori S, Barosi G, et al., Evidence- and consensus-based practice guidelines for the therapy of primary myelodysplastic syndromes. A statement from the Italian Society of Hematology, *Haematologica*, 2002;87:1286–1306.
- Bowen D, Culligan D, Jowitt S, et al., Guidelines for the diagnosis and therapy of adult myelodysplastic syndromes, *Br J Haematol*, 2003;120:187–200.
- Gattermann N, Porter JB, Lopes LF, Seymour J, Consensus statement on iron overload in myelodysplastic syndromes, *Hematol Oncol Clin North Am*, 2005;19(Suppl. 1):18–25.
- Bennett J, Consensus statement on iron overload in myelodysplastic syndromes, *Am J Hematol*, 2008;83(11): 858–61.
- National Comprehensive Cancer Network, NCCN Clinical Practice Guidelines, Myelodysplastic Syndromes, v.2, 2008. Available at: www.nccn.org
- Gattermann N, Clinical consequences of iron overload in myelodysplastic syndromes and treatment with chelators, *Hematol Oncol Clin North Am*, 2005;19(Suppl. 1):13–17.