

Treatment Options in *BRAF*-mutant Metastatic Colorectal Cancer

Marco Cefali,¹ Maria Celeste Palmarocchi¹ and Sara De Dosso^{1,2}

1. Oncology Institute of Southern Switzerland, EOC, Bellinzona, Switzerland; 2. Università della Svizzera Italiana, Lugano, Switzerland

DOI: <https://doi.org/10.17925/OHR.2022.18.2.103>

Treatment of *BRAF*-mutant colorectal cancer (CRC) traditionally represents an unmet need, mainly due to its unfavourable prognostic outlook, limited options for targeted treatment and scarce benefit from epithelial growth-factor receptor (*EGFR*) inhibitors. Recently, the development of *BRAF* V600E inhibitors has expanded the therapeutic armamentarium, although exclusive targeting of *BRAF* has proved to be an unsuccessful strategy due to reactivation of the mitogen-activated protein kinase pathway through multiple escape mechanisms. Combination strategies that exploit simultaneous inhibition of *BRAF*, *EGFR* and/or mitogen-activated protein/extracellular signal-regulated kinase have achieved greater success, with the BEACON CRC trial providing the first evidence for an improvement in survival with a chemotherapy-free approach in pre-treated patients with CRC, leading to regulatory approval for the combination of encorafenib and cetuximab. Subsequent research efforts attempt to build on these foundations, exploring targeted maintenance strategies and conceivably moving the combination towards the first line of therapy soon, as well as laying the foundation for the use of liquid biopsy as a guidance tool in a precision oncology approach.

Keywords

Colorectal cancer, *BRAF* mutation, encorafenib, cetuximab, *EGFR*, targeted therapy

Disclosure: Sara De Dosso discloses advisory board participation, honoraria and travel support from: Amgen, BMS, Ipsen, Merck, MSD, Novartis, Pfizer, Pierre Fabre, Roche and Servier. Marco Cefali and Maria Celeste Palmarocchi have no financial and non-financial conflicts of interest to declare in relation to this paper.

Review process: Double-blind peer review.

Compliance with ethics: This study involves a review of the literature, and did not involve any studies with human or animal subjects performed by any of the authors.

Data availability: Data sharing is not applicable to this article as no datasets were generated or analysed during the writing of this article.

Authorship: The named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval for the version to be published.

Access: This article is freely accessible at [touchONCOLOGY.com](https://touchoncology.com). © Touch Medical Media 2022

Received: 28 February 2022

Accepted: 28 April 2022

Published online: 22 July 2022

Citation: *touchREVIEWS in Oncology & Haematology*. 2022;18(2):103–6

Corresponding author: PD Dr Med Sara De Dosso, Oncology Institute of Southern Switzerland, Via Gallino 12, 6500 Bellinzona, Switzerland. E: sara.dedosso@eoc.ch

Support: No funding was received for the publication of this article.

Until recently, treatment options for metastatic colorectal cancer (CRC) have been limited to a cytotoxic chemotherapy with or without an anti-epithelial growth-factor receptor (*EGFR*) monoclonal antibody, namely cetuximab and panitumumab, or an anti-angiogenic agent such as bevacizumab, aflibercept or ramucirumab. For subsequent lines of treatment, the tyrosine kinase inhibitor regorafenib is an important part of the therapeutic armamentarium.¹ Targeted treatment in metastatic CRC has traditionally been limited to the use of anti-*EGFR* agents in association with chemotherapy in rat sarcoma virus (*RAS*) wild-type tumours.

Despite their target being located on the cell surface, the clinical activity of anti-*EGFR* antibodies does not correlate with *EGFR* membrane expression; instead, alterations in its downstream signalling pathway, most notably in the *RAS* genes, can bypass *EGFR* inhibition and negate the therapeutic benefit.^{2,3} For this reason, confirmation of pan-*RAS* wild-type status is required before starting treatment with an anti-*EGFR* agent. Tumour sidedness also appears to play a pivotal role, with right-sided tumours reporting scarce benefit from anti-*EGFR* agents, despite wild-type *RAS* status.⁴

Recent studies have enhanced knowledge of the molecular complexity of CRC, possibly improving the stratification and clinical management of patients with CRC. Guinney et al. reported a consensus molecular classification for CRC based on gene-expression profiling, identifying four main subtypes: microsatellite-instability (MSI) immune, canonical, metabolic and mesenchymal.⁵ Interestingly, right-sided tumours are more frequently classified as MSI immune or metabolic subtypes. While these consensus molecular subtypes are not currently evaluated in clinical practice, nor routinely employed to guide treatment choice, the identification of specific molecular alterations have started to pave the way to future possibilities for personalized treatment. In particular, personalized treatment strategies are being explored in clinical trials for a variety of molecular alterations: *HER2* amplification (HERACLES-A, MyPathway), *KRAS* G12C mutation (CodeBreaK100, KRYSTAL-1), *NTRK* gene fusions (NAVIGATE), MSI (KEYNOTE-177) and, relevant to the scope of our review, *BRAF* V600E mutation (BEACON CRC, ANCHOR CRC, BREAKWATER, MODUL).^{6–15}

The purpose of this review is to better delineate the novel treatment options in *BRAF*-mutant metastatic CRC.

BRAF mutations in colorectal cancer

The rapidly accelerated fibrosarcoma (RAF) protein family comprises three main isoforms (*ARAF*, *BRAF* and *CRAF*), with *BRAF* being the most frequently mutated.¹⁶ Moreover, *BRAF* is involved in the *RAS*/*RAF*/*MEK*/extracellular signal-regulated kinases (*ERK*) signalling pathway, also known as the mitogen-activated protein kinase (*MAPK*) cascade, driving cell survival, proliferation and differentiation.¹⁶ Mutations in *BRAF* are present in approximately 10% of patients with metastatic

CRC, and are associated with right-sided tumours, age (>60 years), gender (female), high grade of anaplasia, and the presence of lymph nodal and peritoneal metastases.¹⁷ The most frequent alteration observed is the V600E kinase-activating mutation, representing approximately 90% of cases.¹⁸ Generally, *BRAF* mutations are associated with a worse prognosis in CRC, as shown by Venderbosch et al. in a pooled analysis of data from the CAIRO, CAIRO2, COIN and FOCUS studies.¹⁹ Other less frequent mutations, such as *BRAF* D594G or G596N, are known to be prognostically favourable in CRC, resulting in reduced kinase activity.²⁰ *BRAF* mutations are also closely associated with sporadic MSI and can be detected in 40–60% of MSI-high CRC cases; this overlap can be explained by prevalence of the CpG island methylator phenotype (CIMP) in *BRAF*-mutant cancers, leading to hypermethylation of the *MLH1* promoter, and thus to a microsatellite-unstable phenotype.²¹ On the other hand, no *BRAF*-mutated cases have been reported with MSI associated with Lynch syndrome.¹⁷

Interestingly, the unfavourable prognostic effect of *BRAF* mutations appears to be mitigated, not deleted, by MSI, as shown by Manthravadi et al. in a recent meta-analysis that concentrated on the non-metastatic setting.²²

BRAF mutations have been reported as mutually exclusive with *RAS* mutations, and this could be suggestive of a similarly detrimental effect on the efficacy of *EGFR*-directed therapies through downstream activation of the pathway. It has, however, been suggested that mutations other than V600E may not show the same activation behaviour, but no robust data have been provided to confirm this issue.^{23,24}

Current treatment landscape for *BRAF* V600E-mutant colorectal cancer

In 2016, the TRIBE study confirmed FOLFOXIRI (fluorouracil, leucovorin, oxaliplatin and irinotecan) + bevacizumab as an highly effective first line treatment first-line treatment of metastatic CRC, based on a median overall survival (OS) benefit (median OS 19.0 months with FOLFOXIRI + bevacizumab versus 10.7 months with FOLFIRI [folinic acid, fluorouracil and irinotecan] + bevacizumab).²⁵ However, in the same trial, a subgroup analysis failed to identify benefits in the *BRAF*-mutation subgroup of patients. An individual patient data meta-analysis appeared to confirm that the greatest advantage from the triplet + bevacizumab combination was observed in patients with the *KRAS* mutation (who are ineligible for *EGFR*-directed therapies), while those carrying the *BRAF* mutation did not present significant benefits.²⁶

The aim of the phase II FIRE-4.5 study was to determine if cetuximab or bevacizumab were the best drugs to be combined with FOLFOXIRI in patients with *BRAF* V600E-mutated tumours, a population in which first-line prospective data were lacking.²⁷ Based on a review of the literature, the use of *EGFR*-directed therapies in this population is controversial, and the trial officially turned out negative, only detecting a statistically significant progression-free survival (PFS) advantage with bevacizumab versus cetuximab.²⁷ Thus, conventional chemotherapy associated with biologic agents appears to be globally unsatisfactory for this subgroup of patients.

Attempts to improve the dismal prognosis of patients with *BRAF*-mutated metastatic CRC were then concentrated on precision oncology. The development of *BRAF* inhibitors specifically targeted at the V600E mutation appeared to identify a strategy to target the *MAPK* cascade in these patients. Nevertheless, *BRAF* inhibition alone was observed to remove inhibitory feedback on *EGFR* expression,

thus leading to increased *EGFR* density on the cell surface. Thus, shunted signal transduction through CRAF or other associated signalling pathways deleted the effect of the blockade.^{28,29} For these reasons, a rationale for multiple inhibition emerged. For instance, the phosphatidylinositol-3-kinase (*PI3K*)/mammalian target of rapamycin (mTOR) signalling pathway is known to be constitutively activated in a significant portion of patients with CRC (30%) and also, specifically, in *BRAF*-mutant CRC, thus emerging as a possible resistance mechanism to *BRAF*-directed monotherapy.^{30,31} Based on these notions, Coffee et al. explored the efficacy of combined *BRAF* and *PI3K*/mTOR inhibition in genetically engineered mouse models, with interesting signals of activity.³¹ Similarly, preclinical data reported by Prahallad et al. have shown that a combination of *BRAF* and *EGFR* inhibitors seems to produce a synergistic effect in CRC xenografts.²⁹

Clinical data on triple targeting of *PI3K*, *BRAF* and *EGFR* come from van Geel et al., who tested the combination of encorafenib and cetuximab with or without the *PI3K* catalytic subunit α (*PI3KCA*) inhibitor alpelisib in the setting of a phase Ib dose-escalation study for metastatic *BRAF* mutant CRC, with or without *PI3KCA* alterations.³³ Results showed promising clinical activity with an overall response rate (ORR) of 18% and 19% and a median PFS of 3.7 months and 4.2 months, with the doublet and triplet combination, respectively, and an acceptable toxicity.³³

In a subsequent phase II trial, 102 patients were randomized to either encorafenib + cetuximab or encorafenib + cetuximab + alpelisib, evaluating PFS as the primary endpoint.³⁴ While the addition of alpelisib appeared to increase toxicity, the clinical activity was promising, with a mean PFS of 4.2 months with the doublet and 5.4 months with the triplet. OS was evaluated as a secondary endpoint in an interim analysis, resulting in a median OS of 15.2 months with the triplet versus not reached with the doublet.³⁴ More data are required on this treatment approach.

Likewise, the combined inhibition of *BRAF* and *MEK* has a solid preclinical rationale, as evaluated by Corcoran et al.³⁵ For this reason, chemotherapy-free doublet or triplet regimens associating a *BRAF* inhibitor with a *MEK* inhibitor, an anti-*EGFR* antibody or both were investigated. Hyman et al. evaluated this approach in a phase II basket trial, where 37 heavily pre-treated patients with CRC received the *BRAF* inhibitor vemurafenib either alone or in association with cetuximab, resulting in an ORR of 0% and 4%, and disease stability in 50% and 69%, respectively; median PFS in the combination group was 3.7 months and median OS was 7.1 months.³⁶ Similarly, a pilot trial by Yaeger et al. in 15 patients tested the combination of vemurafenib and panitumumab in pre-treated metastatic CRC, with an acceptable safety profile and modest clinical activity; tumour regression was observed in 10 out of 12 patients, but a formal partial response was only defined in two patients, with two more patients obtaining disease stability lasting more than 6 months.³⁷

Doublet therapy with the *BRAF* inhibitor dabrafenib and the *MEK* inhibitor trametinib was tested by Corcoran et al. in 43 patients with *BRAF*-mutant CRC, obtaining interesting signals of activity, with an ORR of 12%, including one complete response, and disease stability in 56% of the study population.³⁸

Based on these data, a new trial explored *BRAF* and *EGFR* inhibition with the doublet of dabrafenib + panitumumab, the triplet of dabrafenib + trametinib + panitumumab, or an alternating doublet of trametinib + panitumumab, in 142 patients with *BRAF* V600E-mutant CRC.³⁹ The

ORR was 0% for trametinib + panitumumab, 10% for dabrafenib + panitumumab and 21% for the triplet, although this greater activity was also associated with increased toxicity. Median PFS was 2.6 months (95% confidence interval [CI], 1.4–2.8 months) with trametinib + panitumumab, 3.5 months (95% CI, 2.8–5.8 months) with dabrafenib + panitumumab, and 4.2 months (95% CI, 4.0–5.6 months) with the triplet.³⁹ The BEACON CRC phase III trial assessed the encorafenib + binimetinib + cetuximab triplet regimen and the encorafenib + cetuximab doublet regimen versus the investigators' choice of irinotecan + cetuximab or FOLFIRI + cetuximab, in 665 patients with *BRAF* V600E-mutated metastatic CRC progressing after first- or second-line treatment and who had not received anti-*EGFR*, anti-*BRAF* or anti-*MEK* agents.¹² Primary endpoints were OS and ORR in the triplet group versus the control group. The secondary endpoint of OS in the doublet group versus the control group was also observed. The median OS was 9.3 months (95% CI, 8.2–10.8 months) for the triplet group and 5.9 months (95% CI, 5.1–7.1) for the control group (hazard ratio [HR] versus control, 0.60 [95% CI, 0.47–0.75]). Median OS for the doublet group was 9.3 months (95% CI, 8.0–11.3 months; HR versus control, 0.61 [95% CI, 0.48–0.77]). Confirmed ORR was 26.8% (95% CI, 21.1–33.1%) for the triplet group, 19.5% (95% CI, 14.5–25.4%) for the doublet group and 1.8% (95% CI, 0.5–4.6%) for the control group. The adverse-event profile was favourable, with grade 3 or higher adverse events in 65.8%, 57.4% and 64.2% of patients in the triplet, doublet and control groups, respectively, and a discontinuation rate due to an adverse event of 9%, 9% and 11%, respectively. One fatal event in the triplet arm and two in the control arm were possibly treatment related.¹²

In summary, the combination of encorafenib and cetuximab improved survival and tumour response with a safety profile not significantly different from standard treatment; however, the addition of binimetinib did not show any benefits in terms of OS and was associated with added toxicity. Patients with high C-reactive protein levels, an Eastern Cooperative Oncology Group performance status of 1, an incomplete primary tumour resection, and metastatic involvement of three or more organs may be candidates for greater benefit from the triplet combination.¹²

Interestingly, an additional analysis demonstrated that patients treated with the double combination showed similar OS regardless of prior therapies (including bevacizumab) or the duration of prior therapy used.⁴⁰

The BEACON CRC trial thus provided the first evidence of improvement in survival from a chemotherapy-free targeted therapy in a prospectively biomarker-defined population of patients with metastatic CRC. This led to regulatory approval by the European Medicines Agency (EMA) for the combination of encorafenib + cetuximab in pre-treated, *BRAF* V600E-mutated metastatic CRC in June 2020.¹²

Preliminary results from the phase II ANCHOR CRC trial were recently presented as an abstract at the 2021 European Society for Medical Oncology World Congress on Gastrointestinal Cancer.¹³ The aim of the trial was to translate the encouraging data from BEACON CRC into the first-line setting, evaluating the combination of encorafenib + binimetinib + cetuximab in a treatment-naïve population of 95 patients with *BRAF* V600E-mutated CRC. Preliminary results showed an ORR of 47.8% and a median OS of 17.2 months, with grade 3 or higher adverse events in 69.5% of patients, thus confirming the activity of the combination in the absence of new safety signals.¹³

The BREAKWATER phase III study is a further step towards first-line use of targeted drugs in this setting.¹⁴ The trial has a safety lead-in phase, enrolling patients with *BRAF* V600E-mutated metastatic CRC who have received no more than one line of treatment; this is followed by a phase III trial evaluating three separate arms in a fully treatment-naïve population. Arm A will receive therapy with encorafenib + cetuximab; arm B will receive the doublet plus standard chemotherapy with FOLFOX (folinic acid, fluorouracil, and oxaliplatin) or FOLFIRI; and the control arm will receive standard chemotherapy alone according to the physician's choice, with or without an anti-VEGF antibody. The primary endpoint is PFS in arm A and arm B versus the control arm, respectively; OS will also be evaluated as a key secondary endpoint.¹⁴

The phase II MODUL trial evaluated novel agents as biomarker-driven maintenance therapy in patients with metastatic CRC treated with FOLFOX + bevacizumab in the first line.¹⁵ It comprised four experimental treatment cohorts and a control cohort employing fluoropyrimidine + bevacizumab. The *BRAF*-mutated cohort received treatment with vemurafenib + cetuximab + 5-fluorouracil and leucovorin; the primary endpoint was PFS. At a median follow-up of 16.4 months, median PFS was 11.6 months in the control arm and 10.0 months in the experimental arm; ORR was 25% versus 50%, respectively.⁴¹ The trial was closed prematurely due to events unrelated to the *BRAF*-mutated cohort, so the results are therefore purely descriptive; however, an exploratory genomic analysis carried out on sequential liquid biopsies identified mutations in the *MAPK* pathway as possible drivers of acquired resistance to the experimental treatment.¹⁵

Future directions

The treatment landscape of *BRAF*-mutant CRC is rapidly changing towards treatments that have demonstrated improved outcomes in this prognostically unfavourable population.^{12,13} Moreover, on-going trials will also enhance comprehension and data on the efficacy of the new target therapy combinations in the first-line setting. Therapeutic intervention on multiple blockades along the *MAPK* pathway is efficient in different experiences; however, pharmacological options for targeting the specific proteins along the signalling cascade are evolving, and selecting the best targets and agents to optimize efficacy and minimize toxicity will be an important issue to be considered for future research.^{12,29,32–39}

Molecular profiling is making a difference: a growing knowledge of the molecular landscape of these tumours, which were traditionally handled according to a 'one size fits all' approach, is making a personalized approach more feasible. Additionally, unravelling the interplay of clinical, immunological and molecular features will become fundamental for both stratification of patients and treatment choices. In this perspective, MSI and *BRAF* V600E-mutant CRC is particularly relevant: on one hand, the exact prognostic outlook of this entity is yet to be clarified, on the other hand, *BRAF* mutations do not seem to hinder response to immunotherapy, which opens the way to a possible sequencing strategy.¹¹

The landscape of clinical and molecular parameters and their evolution over the course of the disease's history is fundamental, as it allows clinicians to monitor tumour progression and resistance through non-invasive approaches. Despite the inconclusive results of the MODUL trial, its strategy of longitudinal monitoring through liquid biopsy is likely to be a part of future investigation methods.¹⁵ The inclusion of patients in clinical trials and the incorporation of these techniques in prospective studies must be a priority to better understand and defeat this illness. □

- Van Cutsem E, Cervantes A, Nordlinger B, et al. Metastatic colorectal cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2014;25:iii–iii9.
- Hecht JR, Mitchell E, Neubauer MA, et al. Lack of correlation between epidermal growth factor receptor status and response to panitumumab monotherapy in metastatic colorectal cancer. *Clin Cancer Res*. 2010;16:2205–13.
- Sorich MJ, Wiese MD, Rowland A, et al. Extended RAS mutations and anti-EGFR monoclonal antibody survival benefit in metastatic colorectal cancer: A meta-analysis of randomized, controlled trials. *Ann Oncol*. 2015;26:13–21.
- Snyder M, Bottiglieri S, Almhanna K. Impact of primary tumor location on first-line bevacizumab or cetuximab in metastatic colorectal cancer. *Rev Recent Clin Trials*. 2018;13:139–49.
- Guinney J, Dienstmann R, Wang X, et al. The consensus molecular subtypes of colorectal cancer. *Nat Med*. 2015;21:1350–6.
- Siena S, Sartore-Bianchi A, Trusolino L, et al. D01* - Final results of the HERACLES trial in HER2 amplified colorectal cancer. *Ann Oncol*. 2016;27:iv39.
- Meric-Bernstam F, Hurwitz H, Raghav KPS, et al. Pertuzumab plus trastuzumab for HER2-amplified metastatic colorectal cancer (MyPathway): an updated report from a multicentre, open-label, phase 2a, multiple basket study. *Lancet Oncol*. 2019;20:518–30.
- Hong DS, Fakih MG, Strickler JH, et al. KRAS G12C inhibition with sotorasib in advanced solid tumors. *N Engl J Med*. 2020;383:1207–17.
- Johnson ML, Ou SHI, Barve M, et al. KRYSTAL-1: Activity and safety of adagrasib (MRTX849) in patients with colorectal cancer (CRC) and other solid tumors harboring a KRAS G12C mutation. *Eur J Cancer*. 2020;138:S2.
- Berlin J, Hong DS, Deeken JF, et al. Efficacy and safety of larotrectinib in patients with TRK fusion gastrointestinal cancer. *J Clin Oncol*. 2020;38:824.
- Shiu K-K, Andre T, Kim TW, et al. KEYNOTE-177: Phase III randomized study of pembrolizumab versus chemotherapy for microsatellite instability-high advanced colorectal cancer. *J Clin Oncol*. 2021;39:6.
- Kopetz S, Grothey A, Van Cutsem E, et al. Encorafenib plus cetuximab with or without binimetinib for BRAF V600E metastatic colorectal cancer: Updated survival results from a randomized, three-arm, phase III study versus choice of either irinotecan or FOLFIRI plus cetuximab (BEACON CRC). *J Clin Oncol*. 2020;38:4001.
- Grothey A, Tabernero J, Taieb J, et al. LBA-5 ANCHOR CRC: a single-arm, phase 2 study of encorafenib, binimetinib plus cetuximab in previously untreated BRAF V600E-mutant metastatic colorectal cancer. *Ann Oncol*. 2020;31:S242–3.
- Kopetz S, Grothey A, Yaeger R, et al. BREAKWATER: randomized phase 3 study of encorafenib (enco) + cetuximab (cetux) ± chemotherapy for first-line (1L) treatment (tx) of BRAF V600E-mutant (BRAF V600E) metastatic colorectal cancer (mCRC). *J Clin Oncol*. 2021;39:TPS3619.
- Schmoll H-J, Arnold D, de Gramont A, et al. MODUL—a multicenter randomized clinical trial of biomarker-driven maintenance therapy following first-line standard induction treatment of metastatic colorectal cancer: an adaptable signal-seeking approach. *J Cancer Res Clin Oncol*. 2018;144:1197–204.
- Markowitz SD, Bertagnolli MM. Molecular origins of cancer: Molecular basis of colorectal cancer. *N Engl J Med*. 2009;361:2449–60.
- Molina-Cerrillo J, San Román M, Pozas J, et al. BRAF mutated colorectal cancer: New treatment approaches. *Cancers*. 2020;12:1571.
- Clarke CN, Kopetz ES. BRAF mutant colorectal cancer as a distinct subset of colorectal cancer: Clinical characteristics, clinical behavior, and response to targeted therapies. *J Gastrointest Oncol*. 2015;6:660–7.
- Venderbosch S, Nagtegaal ID, Maughan TS, et al. Mismatch repair status and BRAF mutation status in metastatic colorectal cancer patients: A pooled analysis of the CAIRO, CAIRO2, COIN, and FOCUS studies. *Clin Cancer Res*. 2014;20:5322–30.
- Cremolini C, Di Bartolomeo M, Amatu A, et al. BRAF codons 594 and 596 mutations identify a new molecular subtype of metastatic colorectal cancer at favorable prognosis. *Ann Oncol*. 2015;26:2092–7.
- Weisenberger DJ, Siegmund KD, Campan M, et al. CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. *Nat Genet*. 2006;38:787–93.
- Manthavadi S, Sun W, Saeed A. Prognostic impact of BRAF V600E mutation in patients with non-metastatic colorectal cancer with microsatellite instability: A systematic review and meta-analysis. *J Clin Oncol*. 2018;36:3597.
- De Roock W, Claes B, Bernasconi D, et al. Effects of KRAS, BRAF, NRAS, and PIK3CA mutations on the efficacy of cetuximab plus chemotherapy in chemotherapy-refractory metastatic colorectal cancer: A retrospective consortium analysis. *Lancet Oncol*. 2010;11:753–62.
- Yaeger R, Kotani D, Mondaca S, et al. Response to anti-EGFR therapy in patients with BRAF non-V600-mutant metastatic colorectal cancer. *Clin Cancer Res*. 2019;25:7089–97.
- Cremolini C, Loupakis F, Antoniotti C, et al. FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: Updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. *Lancet Oncol*. 2015;16:1306–15.
- Cremolini C, Antoniotti C, Stein A, et al. Individual patient data meta-analysis of FOLFOXIRI plus bevacizumab versus doublets plus bevacizumab as initial therapy of unresectable metastatic colorectal cancer. *J Clin Oncol*. 2020;38:3314–24.
- Stintzing S, Heinrich K, Tougeron D, et al. Randomized study to investigate FOLFOXIRI plus either bevacizumab or cetuximab as first-line treatment of BRAF V600E-mutant mCRC: The phase-II FIRE-4.5 study (AIO KRK-0116). *J Clin Oncol*. 2021;39:3502.
- Taleb J, Lapeyre-Prost A, Laurent Puig P, et al. Exploring the best treatment options for BRAF-mutant metastatic colon cancer. *Br J Cancer*. 2019;121:434–42.
- Prahallad A, Sun C, Huang S, et al. Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature*. 2012;483:100–3.
- Baba Y, Noshio K, Shima K, et al. Phosphorylated AKT expression is associated with PIK3CA mutation, low stage, and favorable outcome in 717 colorectal cancers. *Cancer*. 2011;117:1399–408.
- Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet*. 2006;7:606–19.
- Coffee EM, Faber AC, Roper J, et al. Concomitant BRAF and PI3K/mTOR blockade is required for effective treatment of BRAF(V600E) colorectal cancer. *Clin Cancer Res*. 2013;19:2688–98.
- van Geel RMJM, Tabernero J, Elez E, et al. A phase Ib dose-escalation study of encorafenib and cetuximab with or without alpelisib in metastatic BRAF-mutant colorectal cancer. *Cancer Discov*. 2017;7:610–9.
- Tabernero J, Geel RV, Guren TK, et al. Phase 2 results: Encorafenib (ENCO) and cetuximab (CETUX) with or without alpelisib (ALP) in patients with advanced BRAF-mutant colorectal cancer (BRAFM CRC). *J Clin Oncol*. 2016;34:3544.
- Corcoran RB, Dias-Santagata D, Bergethson K, et al. BRAF gene amplification can promote acquired resistance to MEK inhibitors in cancer cells harboring the BRAF V600E mutation. *Sci Signal*. 2010;3:ra84.
- Hyman DM, Puzanov I, Subbiah V, et al. Vemurafenib in multiple nonmelanoma cancers with BRAF V600 mutations. *N Engl J Med*. 2015;373:726–36.
- Yaeger R, Cercek A, O'Reilly EM, et al. Pilot trial of combined BRAF and EGFR inhibition in BRAF-mutant metastatic colorectal cancer patients. *Clin Cancer Res*. 2015;21:1313–20.
- Corcoran RB, Atreya CE, Falchook GS, et al. Combined BRAF and MEK inhibition with dabrafenib and trametinib in BRAF V600-mutant colorectal cancer. *J Clin Oncol*. 2015;33:4023–31.
- Corcoran RB, André T, Atreya CE, et al. Combined BRAF, EGFR, and MEK inhibition in patients with BRAF V600E-mutant colorectal cancer. *Cancer Discov*. 2018;8:428–43.
- Kopetz S, Aderka D, Grothey A, et al. Overall survival (OS) with encorafenib (enco) + cetuximab (cetux) in BEACON CRC: effect of prior therapy for BRAF V600E-mutant metastatic colorectal cancer (mCRC). *J Clin Oncol*. 2021;39:3583.
- Ducreux M, Tabernero J, Grothey A, et al. O-9 5-FU/LV + cetuximab + vemurafenib as maintenance therapy for BRAF-mutant (BRAFMut) metastatic colorectal cancer: Efficacy, safety, and exploratory biomarker findings from Cohort 1 of the MODUL trial. *Ann Oncol*. 2021;32(Suppl. 3):S221–2.