

The Evolving Therapeutic Landscape and Role of Enfortumab Vedotin in Advanced Urothelial Carcinoma: A Systematic Review

Rafee Talukder,¹ Dimitrios Makrakis,² Petros Grivas¹ and Ali Raza Khaki³

1. Department of Medicine, Division of Oncology, Fred Hutchinson Cancer Center, University of Washington, Seattle, WA, USA; 2. Department of Medicine, Jacobi Medical Center, Albert Einstein College of Medicine, New York, NY, USA; 3. Department of Medicine, Division of Oncology, Stanford University, Stanford, CA, USA

The therapeutic landscape in advanced urothelial carcinoma (aUC) has been rapidly evolving over the last 6 years. Enfortumab vedotin (EV) is an antibody–drug conjugate that targets Nectin-4, which is widely expressed in UC. EV is approved by the US Food and Drug Administration for patients with aUC refractory to prior therapy. EV is now being investigated in combination with other therapeutic agents, such as immune checkpoint inhibitors, both in first-line and refractory settings in aUC as well as earlier therapy settings. In this review, we summarized the role of EV in the rapidly evolving therapeutic landscape of aUC.

Keywords

Advanced urothelial carcinoma, antibody-drug conjugate, bladder cancer, enfortumab vedotin, immune checkpoint inhibitors, urothelial carcinoma

Disclosures: Petros Grivas has been a consultant for Aadi Bioscience, AstraZeneca, Astellas Pharma, Boston Gene, Bristol Myers Squibb, CG Oncology, Dyania Health, EMD Serono, Exelixis, Fresenius Kabi, G1 Therapeutics, Genentech/Roche, Gilead Sciences, Guardant Health, Infinity Pharmaceuticals, Janssen, Lucence Health, MSD, Mirati Therapeutics, Pfizer, PureTech, QED Therapeutics, Regeneron Pharmaceuticals, Seattle Genetics, Silverback Therapeutics, 4D Pharma PLC and UroGen. His institution has received grants from Bavarian Nordic, Bristol Myers Squibb, Clovis Oncology, Debiopharm, EMD Serono, G1 Therapeutics, Gilead Sciences, GlaxoSmithKline, MSD, Mirati Therapeutics, Pfizer and QED Therapeutics. Ali Raza Khaki has research collaborations with Natera and Tempus Labs; honoraria from MJH Life Sciences and HMP Global; uncompensated advisory board participation with Seagen/Astellas and uncompensated consultation with Janssen. Rafee Talukder and Dimitrios Makrakis have no financial or non-financial relationships or activities to declare in relation to this article.

Review process: Double-blind peer review.

Compliance with ethics: This article involves a review of the literature and did not involve any studies with human or animal subjects performed by any of the authors.

Data availability: Data sharing is not applicable to this article as no datasets were generated or analysed during the writing of this article.

Authorship: The named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval for the version to be published.

Access: This article is freely accessible at [touchONCOLOGY.com](https://touchoncology.com) © Touch Medical Media 2023

Received: 12 August 2022

Accepted: 23 November 2022

Published online: 2 March 2023

Citation: *touchREVIEWS in Oncology & Haematology*. 2023;19(1):Online ahead of journal publication

Corresponding author: Ali Raza Khaki, Department of Medicine, Division of Oncology, Stanford University, 875 Blake Wilbur Dr, Stanford, CA 94305, USA. E: alikhaki@stanford.edu

Support: No funding was received in the publication of this article.

Bladder cancer is one of the most common malignancies in the USA, with an estimated 81,180 new cases and 17,100 deaths in 2022.¹ Advanced urothelial carcinoma (aUC) includes both locally advanced, unresectable and metastatic urothelial cancer. About 5% of all bladder cancer cases present as *de novo* aUC, while almost 25% of patients present with muscle-invasive bladder cancer (MIBC). About 50% of these patients progress to aUC even with intent-to-cure locoregional therapy, for example, neoadjuvant cisplatin-based chemotherapy (in fit patients) followed by radical cystectomy and pelvic lymph node dissection.² The treatment landscape in aUC is rapidly evolving with the advent of novel therapeutic agents and approaches. Enfortumab vedotin (EV) is one of six new medications that have been approved by the US Food and Drug Administration (FDA) for aUC since 2017 and has been shown to have high efficacy with acceptable toxicity in clinical trials and retrospective studies.^{3–7} This article will briefly review the treatment landscape of aUC in both the first-line and salvage settings. We will then focus on EV in the treatment landscape of aUC and, in particular, on the landmark phase III EV-301 trial.⁵

Methodology

A literature search was performed by the authors from March to May 2022 through PubMed/Medline to identify available articles and studies pertaining to the biology and systemic treatment of MIBC and aUC. The search was performed using the following keywords, separately and in different combinations: “bladder cancer”, “antibody drug conjugate”, “checkpoint inhibition”, “metastatic urothelial carcinoma”, “immunotherapy”, “Sacituzumab govitecan”, “Erdafitinib”, “Enfortumab vedotin”, “target therapy” and “trial”. Current abstracts with preliminary data from international conferences, such as the 2022 American Society of Clinical Oncology (ASCO) Genitourinary Cancers Symposium, 2022 ASCO Annual Meeting and the 2022 European Society for Medical Oncology Congress, were reviewed and used for this systematic review. The authors performed the study selection following the recommendations of the Preferred Reporting Items for Systematic Review and Meta-analysis Statement.⁸

First-line treatment options

First-line therapy for aUC includes platinum-based chemotherapy followed by switch maintenance avelumab for those with a response or stable disease after 4–6 cycles of chemotherapy. For patients deemed fit for cisplatin, cisplatin-based combinations, such as gemcitabine with cisplatin or dose-dense methotrexate, vinblastine, doxorubicin, cisplatin (MVAC), are preferred and have demonstrated overall response rates (ORRs) of up to 50–55%, including approximately 10–15% complete responses (CRs).⁹ Initially, cisplatin was considered the standard of care in patients with aUC as early prospective studies showed remissions ranging from 30 to 40%.^{10–12} A phase III trial investigating cisplatin alone compared with MVAC showed superior response rates for MVAC (39%) compared with cisplatin alone (12%) and improved overall survival (OS) with median of 12.5 versus 8.2 months, respectively.¹³ This trial established cisplatin-based combination to be the standard-of-care therapy. However, the conventional MVAC combination was associated with increased treatment-related adverse effects such as neutropenia, neutropenic fever and mucositis.

A subsequent phase III clinical trial investigating the conventional MVAC treatment versus cisplatin/gemcitabine showed similar OS between the two regimens (hazard ratio [HR] 1.04; 95% confidence interval [CI] 0.82–1.32; $p=0.75$), but cisplatin/gemcitabine showed a better safety profile with fewer rates of neutropenia and neutropenic fever.¹⁴ The phase III clinical trial EORTC-30924 also demonstrated decreased toxicity and the ability to deliver twice the dose of cisplatin and doxorubicin with dose-dense MVAC, with improved progression-free survival (PFS) and ORR compared with MVAC.¹⁵

Almost half of the patients with aUC are unfit for cisplatin-containing chemotherapy due to impaired renal function, poor performance status, symptomatic heart failure, severe underlying neuropathy or hearing loss.¹⁶ For these patients, carboplatin with gemcitabine is the preferred first-line regimen, with an ORR of approximately 40%.¹⁷ Despite initial chemosensitivity, historical outcomes after these platinum-based regimens have been poor, with median PFS (mPFS) of only 6–8 months and median OS (mOS) of 9–14 months. Historically, only 15% of patients have experienced 5-year OS, and the prognosis is particularly poor among patients with visceral metastases for whom the 5-year OS rate has been below 10%.¹⁸ In an effort to improve these outcomes, multiple studies have investigated either biomarkers to aid in patient selection or combination therapies,^{19,20} including chemotherapy with immune checkpoint inhibitor (ICI) and ICI combinations; unfortunately, these trials have not shown a significant OS benefit over platinum-based chemotherapy.^{21–24} An approach that has been shown to be very successful is the switch-maintenance immunotherapy. The JAVELIN Bladder 100 phase III trial showed that the use of switch-maintenance avelumab, an anti-programmed death-ligand 1 (PD-L1), plus best supportive care in patients who had a response or stable disease with first-line platinum-based chemotherapy (cisplatin/gemcitabine or carboplatin/gemcitabine) had significantly prolonged OS and PFS compared with best supportive care alone.²⁵

ICIs have also been investigated and approved in the first-line setting for patients who are not eligible to receive platinum. Atezolizumab was FDA approved for aUC in the first-line setting in patients with PD-L1-high tumours (evaluated using the Ventana SP142 assay [Ventana Medical Systems, Inc.; Tucson, AZ, USA]) who were not eligible to receive cisplatin or for those who were not eligible to receive platinum.²⁶ This approval was based initially on the single-arm phase II IMvigor210 cohort one trial, which showed an ORR of 23% with durable responses and favourable toxicity profile.²⁷ The ORR was later confirmed in the IMvigor130 phase III trial (ClinicalTrials.gov Identifier: NCT02807636); however, atezolizumab did not significantly prolong OS versus chemotherapy.²¹ The indication for atezolizumab in the first-line setting was recently withdrawn.²⁸ Pembrolizumab was initially approved in this first-line setting only for patients who were not eligible to receive cisplatin based on the single arm phase II KEYNOTE-052 trial (ClinicalTrials.gov Identifier: NCT02335424), which showed an ORR of 29%, with durable responses and favourable toxicity profile.²⁹ The subsequent KEYNOTE-361 randomized phase III trial (ClinicalTrials.gov identifier: NCT02853305) did not show longer survival with pembrolizumab versus platinum-based chemotherapy, even among those with a PD-L1 combined positive score of ≥ 10 , which led to a revision in the pembrolizumab label by the FDA to limit use to only those who were not eligible for platinum.^{22,30,31} Several other trials investigating ICI combinations have also been investigated and reported in this setting but have not changed practice.^{23,32} However, some phase III clinical trials are on-going, including Checkmate 901 (nivolumab/ipilimumab versus platinum-based chemotherapy [ClinicalTrials.gov Identifier: NCT03036098]), NILE (durvalumab with

platinum-based chemotherapy versus durvalumab/tremelimumab with platinum-based chemotherapy versus platinum-based chemotherapy alone [ClinicalTrials.gov Identifier: NCT03682068]) and EV-302 (EV with pembrolizumab versus platinum-based chemotherapy [ClinicalTrials.gov Identifier: NCT04223856]).^{32–34}

Novel therapies in the salvage setting

Multiple new therapies have been approved in the salvage setting for patients with aUC who have progressed on first-line therapy. ICIs have been extensively investigated in this setting and have shown durable responses after progression on platinum-based chemotherapy. The phase III KEYNOTE-045 trial (ClinicalTrials.gov Identifier: NCT02256436) demonstrated significantly longer OS with pembrolizumab compared with taxane or vinflunine in patients with progression on first-line platinum-based chemotherapy.³⁵ Atezolizumab, durvalumab, nivolumab and avelumab have also been shown to have durable responses with favourable toxicity profiles compared with historical cytotoxic chemotherapy after progression on first-line chemotherapy in single-arm studies.^{36–39} Pembrolizumab, nivolumab and avelumab have on-going FDA-approved indications for treatment after progression on first-line platinum-based chemotherapy, while atezolizumab and durvalumab also had FDA-approved indications in this setting, although subsequent negative trials led to voluntary withdrawal.^{23,40–47} However, with the advent of the switch-maintenance therapy avelumab in those with a response or stable disease on first-line platinum-based chemotherapy, the utility of second-line ICI has diminished, as many patients receive switch-maintenance avelumab and, thus, require other therapies upon progression.

Erdafitinib, a potent tyrosine kinase inhibitor of fibroblast growth factor receptors 1–4 (*FGFR1–4*) involved in signalling pathways that regulate cell proliferation, migration and differentiation, has received accelerated FDA approval in patients who progress on platinum-based chemotherapy and have susceptible *FGFR2* or *FGFR3* alterations (activating mutation or fusion).⁴⁸ Approximately 20% of patients with bladder cancer and 30–50% of those with upper-tract UC may have such alterations.^{49–53} The open-label BLC2001 phase II trial in those with susceptible *FGFR2* or *FGFR3* alterations and documented progression on platinum-based chemotherapy showed an ORR of 40% and an mPFS of 5.5 months with manageable toxicity.⁵⁴ A follow-up phase III study (THOR) is investigating erdafitinib compared with chemotherapy (vinflunine or docetaxel) or pembrolizumab in patients with aUC with certain *FGFR* alterations (*FGFR3* mutations or *FGFR2/3* fusions) in the platinum-refractory setting.⁵⁵ Erdafitinib was also evaluated in combination with cetrelimab (an anti-PD1) in the first-line setting in the phase Ib/II NORSE trial, with early results showing an ORR of 68%.⁵⁶

Sacituzumab govitecan is an antibody–drug conjugate targeting Trop-2, an epithelial cell surface antigen highly expressed in UCs. It has shown an ORR of 27%, an mPFS of 5 months and an mOS of 11 months in heavily pretreated patients who had progressed on platinum-based therapy and ICI based on the phase II TROPHY-U-01 trial (cohort 1) and has received accelerated FDA approval in this setting.⁵⁷ A confirmatory phase III trial (TROPICS-04 study) is now under way.⁵⁸ Cohort 3 of TROPHY-U01, which investigated the combination of sacituzumab govitecan with pembrolizumab in the platinum-refractory, ICI-naïve, second-line setting reported an ORR of 34% and a disease control rate of 61% with manageable toxicity.⁵⁹ There are several reported and on-going trials investigating novel agents, as monotherapy or in combination with ICIs, in patients who have

progressed on first-line platinum-based chemotherapy. Targets under investigation include poly (ADP-ribose) polymerase (PARP), FGFR, human epidermal growth factor receptor 2 and mammalian target of rapamycin.⁶⁰ Below, we review the landmark phase II/III studies investigating novel agents (with or without ICIs) for aUC. The ATLAS trial, an open-label phase II trial investigating rucaparib monotherapy (a PAR inhibitor) in previously treated aUC regardless of homologous recombination deficiency status, did not show activity in unselected patients.⁶¹ The open-label multiarm BISCAY trial investigating second-line combination durvalumab with relevant biomarker-selected targeted therapies in patients with aUC who progressed on platinum-based cohorts had four arms:

1. combination durvalumab with AZD4547 (an FGFR inhibitor) versus AZD4547 alone in patients with *FGFR3* mutations or *FGFR1–3* fusions
2. combination durvalumab with vistusertib (TORC1/2 inhibitor) in patients with RICTOR amplification or *TSC1/2* mutations
3. combination durvalumab plus olaparib (PARP inhibitor) in patients with *DDR* gene alterations
4. durvalumab alone.

Response rates ranged from 9% to 36% across arms and did not meet specified thresholds of target ORR in any of the cohorts.⁶² Other on-going biomarker-driven studies investigating combination novel agents with ICIs include a phase II trial investigating recombinant albumin fusion protein sEphB4-HSA with pembrolizumab for patients with aUC who had progressed on platinum-based chemotherapy (ClinicalTrials.gov identifier: NCT02717156); the trial demonstrated an ORR of 58%, with an mOS of 15 months in those who expressed EphrinB2.⁶³ The multicohort phase II trial of sitravatinib (a tyrosine kinase inhibitor targeting Tyro3, Axl, and Mer receptors, split family receptors and c-Met) in combination with nivolumab in patients with aUC who had progressed on platinum-based chemotherapy showed an ORR of 31% and an mPFS of 4 months.⁶⁴ MORPHEUS-mUC (ClinicalTrials.gov identifier: NCT03869190) is a phase Ib/II umbrella trial evaluating the safety and efficacy of atezolizumab in combination with multiple agents with different targets (including Nectin-4, PARP, CD47, CD38, DPP-4 and IL-6R) in platinum-refractory aUC.⁶⁵

Enfortumab vedotin

Preclinical data and early phase trials

EV is an antibody–drug conjugate comprising a monoclonal antibody targeting Nectin-4, which is covalently linked to the microtubule-disrupting agent monomethyl auristatin E (MMAE). Nectin-4 is a transmembrane protein that belongs to the nectin family of adhesion molecules. It comprises an extracellular domain containing three immunoglobulin-like subdomains, a transmembrane helix and an intracellular region.⁶⁶ Nectins are thought to mediate Ca²⁺-independent cell–cell adhesion via both homophilic and heterophilic trans-interactions at adherens junctions and are involved in cell movement and proliferation.^{67,68} Nectin-4 has been found to be expressed in multiple cancers, particularly urothelial, breast, lung, pancreatic and ovarian cancers.⁶⁹

In preclinical cancer models, moderate-to-strong Nectin-4 staining was observed in 60% of bladder cancer specimens, 53% of breast cancer specimens, 37% of pancreatic cancer specimens and 27% of lung cancer specimens.⁶⁹ EV was shown to bind to cell surface-expressed Nectin-4 with high affinity and to induce cell death *in vitro* in a dose-dependent manner.⁶⁹ Moderate-to-strong Nectin-4 staining was also observed in 34% of patients with upper tract or renal pelvis UC.⁷⁰ The current literature evaluating Nectin-4 expression in variant histology bladder cancer is limited. A recent study investigating Nectin-4 expression in

bladder cancer revealed that there is a heterogeneity of expression in morphological variants of urothelial cancer and non-urothelial histologic types, with 7/10 (70%) squamous cell carcinomas, 3/11 (28%) micropapillary tumours, 4/6 (66%) adenocarcinomas, 2/4 (50%) nested carcinomas, 5/8 (63%) plasmacytoid, 1/10 (10%) sarcomatoid carcinomas and 0/15 (0%) small cell carcinomas expressing Nectin-4.⁷¹

EV-101 (ClinicalTrials.gov identifier: NCT02091999) was a phase I dose-escalation/dose-expansion study in patients with Nectin-4-positive tumours who had progressed on ≥ 1 line of therapy.⁷² The trial included 155 patients in the aUC cohort. Results showed that EV was well tolerated, with an ORR of 43% regardless of previous treatment.⁷²

Based on these results, a follow-up single-arm, two-cohort phase II trial, EV-201 (ClinicalTrials.gov identifier: NCT03219333), was undertaken.³ Patients receiving EV alone (cohort 1) demonstrated an ORR of 44% (12% CR), with a median duration of response of 7.6 months in patients with aUC who had progressed after platinum-based chemotherapy and ICI. The most common treatment-related adverse events were fatigue (50%), peripheral neuropathy (50%), rash (48%; 75% of which were grade ≤ 2) and decreased appetite (44%). Treatment-related hyperglycaemia was seen in 11% of patients.⁴ These results led to the FDA granting accelerated approval to EV in December 2019 for patients with aUC who have received platinum-based chemotherapy and ICI.⁷³

A separate single-arm cohort of EV-201 included patients with aUC who were not eligible for cisplatin and progressed on anti-PD1/L1 therapy.³ In this cohort, the ORR with EV alone was 52% (20% CR rate) with a similar toxicity profile to the one seen in cohort 1. These findings resulted in the expansion of the FDA label in July 2021 (at the same time as the EV regular approval based on the EV-301 trial) to include treatment for patients with aUC, who are ineligible for cisplatin and have received ≥ 1 prior line of therapy.

EV-301

Following the results from EV-201, a confirmatory phase III trial, EV-301 (ClinicalTrials.gov identifier: NCT03474107), was conducted in patients with aUC who had progression on platinum-based chemotherapy and ICI.⁵ Patients were randomized to receive either EV or chemotherapy (docetaxel, paclitaxel or vinflunine as per the investigator's choice). The primary endpoint was OS, and other endpoints included ORR, PFS, toxicity and disease control rate.

Overall, 301 patients were randomized to EV and 307 patients to chemotherapy. EV significantly prolonged OS (median 12.8 versus 8.9 months; HR for death 0.70; 95% CI 0.56–0.89; $p=0.001$) and mPFS (5.5 versus 3.7 months; HR for progression or death 0.62; 95% CI 0.51–0.75; $p<0.001$) in patients with aUC and progression on platinum-based chemotherapy and ICI. ORR was also significantly higher in the EV group compared with the chemotherapy group (40.6% versus 17.9%, respectively; $p<0.001$), as well as the disease control rate (79.0% versus 53.4%, respectively; $p<0.001$). The incidence of treatment-related adverse events was similar between the two groups, with approximately 90% of patients expressing any adverse event and 50% of patients expressing a grade ≥ 3 event in both groups.⁵ Specific treatment-related adverse events associated with EV included peripheral neuropathy (46.3% any grade, 5.1% grade ≥ 3 ; predominantly sensory neuropathy), rash (43.9% any grade, 14.5% grade ≥ 3) and hyperglycaemia (6.4% any grade, 4% grade ≥ 3 , including 0.3% grade 5). Early recognition of toxicity and proper management is critical.⁷⁴ Based on these results, in July 2021, the FDA granted regular approval to EV for patients with aUC after progression

Table 1: Pivotal enfortumab vedotin trial in bladder cancer

	Description	Primary endpoint	Secondary endpoint
EV-101 ⁷²	Phase I dose escalation-/dose-expansion study in patients with Nectin-4-positive tumours, including aUC	Toxicity: Well tolerated. Fatigue (53%), alopecia (46%) and decreased appetite (42%) were the most common TRAEs	ORR 43%; mDOR 7.7 months (95% CI 5.6–9.6)
EV-201 ³	Phase II single-arm, two-cohort study investigating EV in patients with previously treated aUC		
	Cohort 1: Patients received prior platinum-based chemotherapy and ICI ⁴	ORR 44.0% (95% CI 35.1–53.2)	mDOR 7.6 months (range 0.3–24.6); mOS 12.40 months (95% CI 9.46–15.57); mPFS 5.8 months (95% CI 4.9–7.5)
	Cohort 2: Patients with aUC who were not eligible for cisplatin and who had not had prior chemotherapy received ICI ³	ORR 51.6% (95% CI 40.8–62.4)	mDOR 6.00 months (range 0.95–11.30); mOS 14.7 months (95% CI 10.5–18.2); mPFS 5.8 months (95% CI 5.0–8.3)
EV-301 ⁵	Phase III open-label trial investigating EV versus chemotherapy (provider choice) in patients with aUC who had previously received platinum-based chemotherapy and ICI	mOS 12.88 months (95% CI 0.56–0.89)	mPFS 5.55 months (95% CI 0.51–0.75); ORR 40.6% (95% CI 34.9–46.5); Toxicity: ≥G3 TRAE 51.4%
EV-103 ⁷⁷	Phase Ib/II multicohort trial investigating EV alone or in combination in patients with MIBC and aUC		
	Cohort A: Single-arm cohort investigating EV plus P in first-line cisplatin-ineligible aUC ⁷⁷	ORR 73.3% (95% CI 58.1–85.4); Toxicity: neuropathy (56%, 4.4% ≥G3), fatigue (51.1%, 11.1% ≥G3), alopecia (48.9%)	mDOR: 25.6 months (95% CI 8.3–NR); mOS: NR; mPFS: 12.3 months (95% CI 8.0–NR)
	Cohort H: Single-arm cohort investigating neoadjuvant EV in patients with MIBC who are not eligible for cisplatin ⁷⁸	pCR: 36.4%	Pathological downstaging rate: 50%; Toxicity: fatigue (45.5%), alopecia (36.4%), dysgeusia (36.4%)
	Cohort K: Two-arm cohort investigating EV alone or combination EV plus P as first-line treatment for patients with aUC not eligible for cisplatin ⁷⁹	ORR: EV + P: 64.5% (95% CI 52.7–75.1); EV alone: 45.2% (95% CI 33.5–57.3)	DOR: EV + P: NR; EV alone: 13.2 months

CI = confidence interval; EV = enfortumab vedotin; ICI = immune checkpoint inhibitor; MIBC = muscle-invasive bladder cancer; NR = not reached; ORR = objective response rate; P = pembrolizumab; TRAE = treatment-related adverse events; aUC = advanced urothelial carcinoma; mDOR = median duration of response; mOS = median overall survival; mPFS = median progression-free survival; pCR = pathological complete response rate.

on prior platinum-based chemotherapy and ICI.⁷⁵ Longer follow-up data presented at the 2022 annual ASCO meeting confirmed the above data.⁷⁶

Table 1 summarizes the key findings of the pivotal trials investigating EV.^{77–79}

Is there an optimal sequence of treatment in the second line and beyond?

While the results from EV-301 are impressive and show that EV has a superior OS, PFS and ORR compared with single-agent chemotherapy, a remaining question concerns the optimal sequencing of therapies for aUC. EV-301 randomized patients to either EV or investigator-chosen single-agent chemotherapy (taxane or vinflunine).⁵ Outcomes with single-agent chemotherapy in this setting remain poor, with ORR ranging from around 13% to 30%.^{80–82} Further, several new agents have been approved in the salvage setting over the last few years for patients who progressed on first-line platinum-based chemotherapy, including ICI (pembrolizumab, avelumab, nivolumab) for patients who did not receive avelumab maintenance, erdafitinib (for those with susceptible *FGFR2* or *FGFR3* alterations) and sacituzumab govitecan.^{41–43,48,83} However, there are currently no head-to-head trials investigating response and survival with EV compared with these other approved agents. Consequently, the optimal treatment sequence and when to use EV needs to be determined on a patient-by-patient basis. EV is currently approved for patients who are not eligible to receive cisplatin who have progressed on both platinum-based therapy and ICI and for patients who are not eligible for cisplatin who progress on ≥1 line of therapy. For patients treated with platinum-based chemotherapy followed by switch maintenance avelumab, EV would be a very reasonable subsequent therapy after progression on avelumab. The same is also true for those who receive second-line ICI after progression on platinum-based

chemotherapy. However, a retrospective study has suggested that patients with liver and/or bone metastases may have a lower response rate and shorter survival with ICI compared with those without liver and/or bone metastases,⁸⁴ whereas the same may not be the case with EV.⁶ Therefore, in selected patients with high-risk features (e.g. visceral/bone metastasis), EV or erdafitinib (for patients with susceptible *FGFR2* or *FGFR3* alterations) could potentially be considered prior to ICI, in the absence of head-to-head comparisons, while awaiting for the THOR phase III trial and large real-world datasets.^{85,86}

There may also be relevant medical comorbidities that could impact the therapy sequence in aUC. For example, in the EV clinical trials, including EV-301, patients with (sensory or motor) neuropathy grade ≥2, uncontrolled diabetes mellitus (defined as glycated haemoglobin ≥8 or between 7 and <8 with symptoms, such as polyuria and polydipsia), active keratitis or corneal ulcerations were excluded, given the concern that treatment-related adverse events could worsen baseline comorbidity. Therefore, if a patient has any of these medical comorbidities, alternative therapies (e.g. sacituzumab govitecan) or enrolling in a clinical trial (e.g. TROPICS-04) could be considered.⁵⁸ In addition, in a patient with an *FGFR2*- or *FGFR3*-activating mutation or fusion, erdafitinib could be considered prior to or after EV, given the lack of head-to-head comparison between erdafitinib and EV. Although no prospective data are available, a small retrospective study showed that, in patients with *FGFR3* alteration, receiving EV prior to erdafitinib was associated with shorter PFS and lower ORR compared with receiving EV after the progression on erdafitinib.⁸⁷ Another small study showed that none of seven patients with *FGFR3* alteration responded to FGFR inhibitor given post EV, while one patient who received FGFR inhibitor prior to EV responded.⁸⁸ However, larger and prospective studies are needed to validate and inform optimal therapy sequence in aUC.

Use of biomarkers to evaluate response to enfortumab vedotin

Another potential way to further guide therapy selection would be to identify predictive biomarkers. Predictive biomarkers could help identify which patients are likely to benefit more from EV and could further inform treatment decisions, including optimal sequence of therapies.

Nectin-4 expression remains an elusive biomarker, and on-going studies are investigating its efficacy as a reliable predictive biomarker.^{34,89–91} There seems to be variation between different studies investigating Nectin-4 expression in UC. Nectin-4 expression in tissue is calculated and quantified using the H-score, a histochemical scoring system determined by the staining intensity of the cells (ranging from 0 for no signal to 3 for strong signal) multiplied by the percentage (%) of cells stained at a given intensity. H-scores range from 0 to 300, with a score of 0–99 defined as negative or weak, 100–99 as moderate and a score of 200–300 as strong. In a study of 524 primary bladder tumours, moderate-to-strong H-scores (H-score ≥ 100) were found in only 63% of samples.⁶⁹ However, EV-101 and EV-201 revealed that Nectin-4 expression was observed in the vast majority of patients. In the phase I trial EV-101, Nectin-4 expression was initially a requirement for study enrolment; however, almost all of the screened samples exhibited high levels of Nectin-4 expression by immunohistochemistry (IHC), and thus, the protocol was amended, and the requirement was removed.⁷² Consequently, Nectin-4 expression was also not required for entry in either EV-201 or EV-301. In the EV-201 trial, all 120 patients had detectable Nectin-4 expression with a median H-score of 290.⁴ Tissue samples were obtained for an exploratory analysis of Nectin-4 expression in EV-301, and results are still pending.⁵ Possible discrepancies in Nectin-4 staining between EV trials and other studies investigating Nectin-4 expression could be due to the different antibodies used in the IHC testing. Given this variability, IHC expression of Nectin-4 might not be a reliable predictive biomarker for EV response. However, the quantification of Nectin-4 through other methods, such as Nectin-4 mRNA expression (using techniques such as RNAseq or Nanostring transcription), are currently under investigation as potential biomarkers. For example, a retrospective cohort study investigating Nectin-4 mRNA expression in patients with localized bladder tumours across different molecular subtypes found that mRNA expression was heterogeneous across the different subtypes and significantly increased in luminal subtypes.⁹² Further, *in vitro* analysis revealed that decreased expression of Nectin-4 in luminal cells led to resistance to EV.⁹² This may suggest that Nectin-4 RNA and molecular subtyping might be potential biomarkers, though further prospective validation is needed.

Genomic sequencing could also provide insights into potential molecular biomarkers. In a retrospective analysis of next-generation sequencing data of patients with aUC treated with EV, the presence of *TP53* and the absence of *CDKN2A* and *CDKN2B* alterations were associated with favourable response and improved outcomes, suggesting that these might be potential biomarkers that might predict a response to EV but require further validation.⁹³

Clinical biomarkers may also play a pivotal role in helping inform treatment decisions. The Bellmunt risk factors (Eastern Cooperative Oncology Group performance status < 0 , liver metastasis, haemoglobin < 10 g/dL) were developed based on salvage chemotherapy trials and have been used for clinical prognostication and clinical trial benchmarking.⁹⁴ In addition, a retrospective study using a large, multinational database of patients with aUC treated with ICI showed that, among patients with aUC treated with second-line ICI after progression after first-line platinum-based chemotherapy, time to second-line ICI < 3 months after initiation

of first-line chemotherapy was associated with worse outcomes to second-line ICI compared with those who initiated second-line ICI < 6 months after initiation of first-line chemotherapy.⁹⁵ In a separate study using the same database, patients with bone and/or liver metastases in aUC treated with ICI were associated with worse outcomes compared with those without bone and/or liver metastases.⁸⁴ Perhaps, early progression on previous treatment lines, such as platinum-based chemotherapy and/or ICI, as well as the site of metastases, could also be prognostic biomarkers; however, a predictive role, particularly for EV versus other therapy, remains questionable. Large-scale retrospective and prospective analyses are needed to further investigate and validate both clinical and molecular biomarkers for therapy.

Another useful way translational studies may inform our understanding of EV is through the elucidation of mechanisms of resistance. Putative mechanisms of resistance to EV have not yet been fully understood. In preclinical mouse models with breast cancer, tumours resistant to an EV-like antibody–drug conjugate, comprising a human anti-Nectin-4 monoclonal antibody conjugated to MMAE, named N41mab-vcMMAE, showed increased expression of *ABCB1*, which encodes the multidrug resistance protein MDR-1/P-glycoprotein.⁹⁶ The overexpression of this protein has been associated with chemotherapy resistance of tumour cells by preventing the intracellular accumulation of cytotoxic agents.⁹⁷ Interestingly, the preclinical model showed that the addition of P-glycoprotein pharmacological inhibitors restored the sensitivity of resistant tumour cells to N41mab-vcMMAE.⁹⁶ Preclinical models have also shown that loss of Nectin-4 expression was also associated with EV resistance.⁹² On-going studies and biomarker analyses are needed to help further understand mechanisms of resistance to EV and other antibody–drug conjugates.

Looking ahead to the future of enfortumab vedotin: earlier therapy settings?

Given the favourable results of EV in the salvage setting for aUC, considerable enthusiasm remains for exploring whether EV can be moved to earlier therapy lines, either alone or in combination with other therapies. Both the EV-103 and EV-302 trials are investigating response and outcomes with first-line EV alone or in combination with pembrolizumab. EV-103 is a phase Ib/II multicohort, dose-escalation and dose-expansion trial investigating the safety and activity of EV in combination with other agents commonly used in aUC, such as pembrolizumab or cisplatin.⁷⁷ EV-103 cohort A assessed the safety and activity of a combination of EV and pembrolizumab as first-line therapy in patients with aUC not eligible for cisplatin and showed a manageable safety profile and very promising anti-tumour activity, with a confirmed ORR of 73% (17.8% CR), a median response duration of 25.6 months, an mPFS of 12.3 months, mOS not reached, and a median follow-up time of 24.9 months.⁷⁷ Early results from a follow-up two-arm cohort K of EV plus pembrolizumab (the other treatment arm was EV alone, but the trial was not powered to compare the two arms) as first-line treatment for patients not eligible for cisplatin with aUC showed an ORR of 64.5% in the combination arm and 45.2% with EV alone, with a few differences in the toxicity profiles.⁷⁹

EV-302 is a randomized phase III trial investigating a combination of EV plus pembrolizumab versus gemcitabine plus cisplatin or carboplatin (followed by switch-maintenance avelumab, when appropriate, based on a subsequent trial amendment) in previously untreated patients with aUC eligible to receive platinum.³⁴ The primary endpoints of EV-302 are OS and PFS; this trial has not reported data yet. Given the established role of EV after platinum-based chemotherapy and ICI, it will

Table 2: Ongoing trials investigating enfortumab vedotin in bladder cancer

Trial	Description	Primary endpoint(s)
EV-103 Cohort L ⁸⁹	Single-arm cohort investigating perioperative EV monotherapy in patients with MIBC who are not eligible for cisplatin	pCR
EV-302 ³⁴	Phase III trial investigating combination EV plus pembrolizumab versus gemcitabine plus cisplatin or carboplatin (followed by maintenance avelumab, when appropriate) in previously untreated patients with aUC eligible for platinum	OS, PFS
KEYNOTE-905/EV-303 ⁹¹	Phase III trial investigating perioperative combination EV with pembrolizumab versus no experimental perioperative therapy in patients with MIBC who are not eligible to receive cisplatin	pCR, event-free survival
KEYNOTE-B15/EV-304 ⁹⁰	Phase III trial investigating perioperative EV plus pembrolizumab versus neoadjuvant chemotherapy with cisplatin plus gemcitabine in patients with MIBC eligible for cisplatin	pCR, event-free survival
VOLGA ⁹⁸	Phase III trial investigating the efficacy and safety of neoadjuvant EV plus durvalumab/tremelimumab versus EV plus durvalumab versus no perioperative therapy in patients with MIBC who are not eligible for cisplatin	pCR, event-free survival

EV = enfortumab vedotin; MIBC = muscle-invasive bladder cancer; OS = overall survival; PFS = progression-free survival; aUC = advanced urothelial carcinoma; pCR = pathological complete response.

be important to ensure whenever possible that the trial participants in the control (chemotherapy) arm have access to the appropriate off-trial, post-protocol therapies (e.g. ICI, EV) to best inform whether upfront EV/pembrolizumab truly improves OS in this population. Moreover, it will be important to assess the proportion of patients who receive cisplatin as well as the proportion who receive switch maintenance avelumab in the control chemotherapy arm of the EV-302 trial. The question is whether the FDA will provide accelerated approval to EV/pembrolizumab as upfront therapy in patients with aUC not eligible to receive cisplatin based on the results of the EV-103 trial before the EV-302 phase III trial is completed. A future discussion may also include the evaluation of potential dose de-escalation strategies looking at efficacy, safety, quality of life and cost-effectiveness.

In addition, on-going trials are investigating the role of EV, either as monotherapy or in combination with pembrolizumab or other ICI in the perioperative setting.^{89-91,98} Preclinical data have demonstrated the ability of EV to induce immunogenic cell death and enhance tumour cell killing through the 'bystander effect'.⁹⁹ Preliminary data from EV-103 cohort H, a single-arm cohort investigating neoadjuvant EV in patients with MIBC who are eligible to receive cisplatin, has shown a pathological downstaging rate of 50% and a pathological CR rate of 36.4%.⁷⁸ EV-103 cohort L is investigating perioperative EV monotherapy in patients with MIBC not eligible for cisplatin.⁸⁹ Patients will receive three cycles of neoadjuvant EV followed by radical cystectomy and then six additional cycles of adjuvant EV. The primary endpoint is the pathological CR rate. KEYNOTE-B15/EV-304 (ClinicalTrials.gov identifier: NCT04700124) is a phase III trial investigating perioperative EV plus pembrolizumab versus neoadjuvant chemotherapy with cisplatin/gemcitabine in patients with MIBC who are not eligible to receive cisplatin.⁹⁰ Patients randomized to receive EV plus pembrolizumab

will receive four cycles of neoadjuvant EV plus pembrolizumab followed by five cycles of adjuvant EV and 13 cycles of adjuvant pembrolizumab after radical cystectomy. Primary endpoints are pathological CR rate and event-free survival. KEYNOTE-905/EV-303 (ClinicalTrials.gov identifier: NCT03924895) is a phase III trial investigating a perioperative combination of EV with pembrolizumab versus no experimental perioperative therapy in patients with MIBC who are not eligible for cisplatin.⁹¹ Primary endpoints are pathological CR rate and event-free survival. VOLGA (ClinicalTrials.gov identifier: NCT04960709) is a phase III trial investigating the efficacy and safety of neoadjuvant EV plus durvalumab/tremelimumab versus EV plus durvalumab versus no experimental perioperative therapy in cisplatin-ineligible patients with MIBC.⁹⁸ These trials are actively enrolling patients and have the potential to change the treatment paradigm of MIBC. Many active, on-going trials are investigating combinatorial strategies with EV in both the perioperative and the aUC setting. Table 2 lists examples of on-going trials investigating EV in this cancer.

Conclusion

EV is a highly active drug that has already made a very positive impact on aUC. EV-301 was a landmark phase III trial that led to the regular FDA approval of EV in patients who progress after platinum-based therapy and ICI. EV can also be used as a second-line therapy in patients who are not eligible to receive cisplatin. Optimal patient and provider education, recognition and careful management of toxicity by a multidisciplinary team remain essential.⁷⁴ However, multicentre collaborations, randomized trials with approved agents, promising combinations and biomarker-driven studies can help to determine which patients can benefit more from EV and to elucidate the most appropriate therapy setting and sequence. Moreover, biomarker discovery and validation require translational studies and can inform resistance mechanisms. □

- American Cancer Society. Key statistics for bladder cancer. www.cancer.org/cancer/bladder-cancer/about/key-statistics.html (date last accessed 2 May 2022).
- Alfred Witjes J, Lebre T, Comperat EM, et al. Updated 2016 EAU guidelines on muscle-invasive and metastatic bladder cancer. *Eur Urol*. 2017;71:462-75. DOI: 10.1016/j.eururo.2016.06.020
- Yu EY, Petrylak DP, O'Donnell PH, et al. Enfortumab vedotin after PD-1 or PD-L1 inhibitors in cisplatin-ineligible patients with advanced urothelial carcinoma (EV-201): A multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2021;22:872-882. DOI: 10.1016/S1470-2045(21)00094-2
- Rosenberg JE, O'Donnell PH, Balar AV, et al. Pivotal trial of enfortumab vedotin in urothelial carcinoma after platinum and anti-programmed death 1/programmed death ligand 1 therapy. *J Clin Oncol*. 2019;37:2592-2600. DOI: 10.1200/JCO.19.01140
- Powles T, Rosenberg JE, Sonpavde GP, et al. Enfortumab vedotin in previously treated advanced urothelial carcinoma. *N Engl J Med*. 2021;384:1125-1135. DOI: 10.1056/NEJMoa2035807
- Koshkin VS, Henderson N, James M, et al. Efficacy of enfortumab vedotin in advanced urothelial cancer: Analysis from the Urothelial Cancer Network to Investigate Therapeutic Experiences (UNITE) study. *Cancer*. 2022;128:1194-1205. DOI: 10.1002/cncr.34057
- US Food and Drug Administration. FDA grants regular approval to enfortumab vedotin-efjv for locally advanced or metastatic urothelial cancer. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-regular-approval-enfortumab-vedotin-efjv-locally-advanced-or-metastatic-urothelial-cancer
- Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160.
- Powles T, Bellmunt J, Comperat E, et al. Bladder cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol*. 2022;33:244-258. DOI: 10.1016/j.annonc.2021.11.012
- Merrin C. Treatment of advanced bladder cancer with cis-diamminedichloroplatinum (II NSC 119875): A pilot study. *J Urol*. 1978;119:493-495.
- Yagoda A. Phase II trials with cis-dichlorodiammineplatinum(II) in the treatment of urothelial cancer. *Cancer Treat Rep*. 1979;63:1565-1572.
- Rosoff AH, Talley RW, Stephens R, et al. Phase II evaluation of cis-dichlorodiammineplatinum(II) in advanced malignancies of the genitourinary and gynecologic organs: A Southwest Oncology Group Study. *Cancer Treat Rep*. 1979;63:1557-1564.
- Loehrer PJ, Einhorn LH, Elson PJ, et al. A randomized comparison of cisplatin alone or in combination with methotrexate, vinorelbine, and doxorubicin in patients with metastatic urothelial carcinoma: A cooperative group study. *J Clin Oncol*. 1992;10:1066-1073. DOI: 10.1200/JCO.1992.10.7.1066
- von der Maase H, Hansen SW, Roberts JT, et al. Gemcitabine and cisplatin versus methotrexate, vinorelbine, doxorubicin, and cisplatin in advanced or metastatic bladder cancer: Results of a large, randomized, multinational, multicenter, phase III study. *J Clin Oncol*. 2000;18:3068-3077. DOI: 10.1200/JCO.2000.18.17.3068
- Sternberg CN, de Mulder PH, Schornagel JH, et al. Randomized phase III trial of high-dose-intensity methotrexate, vinorelbine, doxorubicin, and cisplatin (MVAC) chemotherapy and recombinant human granulocyte colony-stimulating factor versus classic MVAC in advanced urothelial tract tumors:

- European Organization for Research and Treatment of Cancer Protocol no. 30924. *J Clin Oncol*. 2001;19:2638–2646. DOI: 10.1200/JCO.2001.19.10.2638
16. Dash A, Galsky MD, Vickers AJ, et al. Impact of renal impairment on eligibility for adjuvant cisplatin-based chemotherapy in patients with urothelial carcinoma of the bladder. *Cancer*. 2006;107:506–513. DOI: 10.1002/cncr.22031
 17. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/carboplatin and methotrexate/carboplatin/vinorelbine in patients with advanced urothelial cancer who are unfit for cisplatin-based chemotherapy: EORTC study 30986. *J Clin Oncol*. 2012;30:191–199. DOI: 10.1200/JCO.2011.37.3571
 18. von der Maase H, Sengelov L, Roberts JT, et al. Long-term survival results of a randomized trial comparing gemcitabine plus cisplatin, with methotrexate, vinorelbine, doxorubicin, plus cisplatin in patients with bladder cancer. *J Clin Oncol*. 2005;23:4602–4608. DOI: 10.1200/JCO.2005.07.757
 19. Rizzo A, Mollica V, Massari F. Expression of programmed cell death ligand 1 as a predictive biomarker in metastatic urothelial carcinoma patients treated with first-line immune checkpoint inhibitors versus chemotherapy: A systematic review and meta-analysis. *Eur Urol Focus*. 2022;8:152–159. DOI: 10.1016/j.euf.2021.01.003
 20. Rizzo A, Mollica V, Santoni M, et al. Impact of clinicopathological features on immune-based combinations for advanced urothelial carcinoma: A meta-analysis. *Future Oncol*. 2022;18:739–748. DOI: 10.2217/fon-2021-0841
 21. Galsky MD, Arjia JA, Bamias A, et al. Atezolizumab with or without chemotherapy in metastatic urothelial cancer (IMvigor130): A multicentre, randomised, placebo-controlled phase 3 trial. *Lancet*. 2020;395:1547–1557. DOI: 10.1016/S0140-6736(20)30230-0
 22. Powles T, Csösz T, Özgüroğlu M, et al. Pembrolizumab alone or combined with chemotherapy versus chemotherapy as first-line therapy for advanced urothelial carcinoma (KEYNOTE-361): A randomised, open-label, phase 3 trial. *Lancet Oncol*. 2021;22:931–945. DOI: 10.1016/S1470-2045(21)00152-2
 23. Powles T, van der Heijden MS, Castellano D, et al. Durvalumab alone and durvalumab plus tremelimumab versus chemotherapy in previously untreated patients with unresectable, locally advanced or metastatic urothelial carcinoma (DANUBE): A randomised, open-label, multicentre, phase 3 trial. *Lancet Oncol*. 2020;21:1574–1588. DOI: 10.1016/S1470-2045(20)30541-6
 24. Galsky MD, Powles T, Li S, et al. A phase 3, open-label, randomized study of nivolumab plus ipilimumab or standard of care (SOC) versus SOC alone in patients (pts) with previously untreated unresectable or metastatic urothelial carcinoma (muc; checkmate 901). *J Clin Oncol*. 2018;36(Suppl. 15):TPS4588–TPS4588.
 25. Powles T, Park SH, Voog E, et al. Avelumab maintenance therapy for advanced or metastatic urothelial carcinoma. *N Engl J Med*. 2020;383:1218–1230. DOI: 10.1056/NEJMoa2002788
 26. US Food and Drug Administration. FDA approves new, targeted treatment for bladder cancer. www.fda.gov/news-events/press-announcements/fda-approves-new-targeted-treatment-bladder-cancer#:~:text=The%20U.S.%20Food%20and%20Drug,treat%20this%20type%20of%20cancer (date last accessed 2 May 2022).
 27. Balar AV, Galsky MD, Rosenberg JE, et al. Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: A single-arm, multicentre, phase 2 trial. *Lancet*. 2017;389:67–76. DOI: 10.1016/S0140-6736(16)32455-2
 28. Genentech. Genentech provides update on tecentriq U.S. indication for previously untreated metastatic bladder cancer. www.gene.com/media/statements/ps_112822 (date last accessed 4 May 2022).
 29. Vuky J, Balar AV, Castellano D, et al. Long-term outcomes in KEYNOTE-052: phase II study investigating first-line pembrolizumab in cisplatin-ineligible patients with locally advanced or metastatic urothelial cancer. *J Clin Oncol*. 2020;38:2658–2666. DOI: 10.1200/JCO.2022.40.6_suppl.432
 30. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (muc). *J Clin Oncol*. 2019;37(Suppl. 7):451–451. DOI: 10.1200/JCO.2019.37.7_suppl.451
 31. Merck. FDA approves updated indication for merck’s KEYTRUDA® (pembrolizumab) for treatment of certain patients with urothelial carcinoma (bladder cancer). www.merck.com/news/fda-approves-updated-indication-for-merck-keytruda-pembrolizumab-for-treatment-of-certain-patients-with-urothelial-carcinoma-bladder-cancer/ (date last accessed 4 May 2022).
 32. Loriot Y, Grivas P, De Wit R, et al. First-line pembrolizumab (pembro) with or without lenvatinib (lenva) in patients with advanced urothelial carcinoma (LEAP-01): A phase 3, randomized, double-blind study. *J Clin Oncol*. 2022;40(Suppl. 6):432. DOI: 10.1200/JCO.2022.40.6_suppl.432
 33. Galsky MD, Necchi A, Sridhar SS, et al. A phase III, randomized, open-label, multicenter, global study of first-line durvalumab plus standard of care (soc) chemotherapy and durvalumab plus tremelimumab, and soc chemotherapy versus soc chemotherapy alone in unresectable locally advanced or metastatic urothelial cancer (NILE). *J Clin Oncol*. 2021;39(Suppl. 6):TPS504–TPS504. DOI: 10.1200/JCO.2021.39.6_suppl.TPS504
 34. Van Der Heijden MS, Gupta S, Galsky MD, et al. Study EV-302: A two-arm, open-label, randomized controlled phase 3 study of enfortumab vedotin in combination with pembrolizumab versus chemotherapy in previously untreated advanced urothelial carcinoma (aUC) (trial in progress). *J Clin Oncol*. 2022;40(Suppl. 6):TPS589–TPS589.
 35. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as second-line therapy for advanced urothelial carcinoma: a phase 2 trial. *N Engl J Med*. 2017;376:1015–26. DOI: 10.1056/NEJMoa1613683
 36. Galsky MD, Sacci A, Szabo PM, et al. Nivolumab in patients with advanced platinum-resistant urothelial carcinoma. Efficacy, safety, and biomarker analyses with extended follow-up from checkmate. *Clin Cancer Res*. 2020;26:5120–5128. DOI: 10.1158/1078-0432.CCR-19-4162
 37. Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): Pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol*. 2018;19:51–64. DOI: 10.1016/S1470-2045(17)30900-2
 38. Rosenberg JE, Hoffman-Censits J, Powles T, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: A single-arm, multicentre, phase 2 trial. *The Lancet*. 2016;387:1909–1920. DOI: 10.1016/S0140-6736(16)00561-4
 39. Powles T, O’Donnell PH, Massard C, et al. Efficacy and safety of durvalumab in locally advanced or metastatic urothelial carcinoma. *JAMA Oncol*. 2017;3:e172411. DOI: 10.1001/jamaoncol.2017.2411
 40. Powles T, Durán J, van der Heijden MS, et al. Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): A multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2018;391:748–757. DOI: 10.1016/S0140-6736(17)33297-X
 41. US Food and Drug Administration. Pembrolizumab (keytruda): Advanced or metastatic urothelial carcinoma. www.fda.gov/drugs/resources-information-approved-drugs/pembrolizumab-keytruda-advanced-or-metastatic-urothelial-carcinoma (date last accessed 2 May 2022).
 42. US Food and Drug Administration. Nivolumab for treatment of urothelial carcinoma. www.fda.gov/drugs/resources-information-approved-drugs/nivolumab-treatment-urothelial-carcinoma (date last accessed 2 May 2022).
 43. US Food and Drug Administration. FDA grants accelerated approval to avelumab for urothelial carcinoma. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-avelumab-urothelial-carcinoma (date last accessed 2 May 2022).
 44. US Food and Drug Administration. Atezolizumab for urothelial carcinoma. www.fda.gov/drugs/resources-information-approved-drugs/atezolizumab-urothelial-carcinoma (date last accessed 2 May 2022).
 45. US Food and Drug Administration. Durvalumab (imfinzi). www.fda.gov/drugs/resources-information-approved-drugs/durvalumab-imfinzi (date last accessed 2 May 2022).
 46. Genentech. Genentech provides update on tecentriq U.S. indication in prior-platinum treated metastatic bladder cancer. www.gene.com/media/press-releases/14898/2021-03-07/genentech-provides-update-on-tecentriq-u (date last accessed 2 May 2022).
 47. AstraZeneca. Voluntary withdrawal of imfinzi indication in advanced bladder cancer in the US. www.astrazeneca.com/media-centre/press-releases/2021/voluntary-withdrawal-imfinzi-us-bladder-indication.html (date last accessed 2 May 2022).
 48. FDA grants accelerated approval to erdafitinib for metastatic urothelial carcinoma. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-erdafitinib-metastatic-urothelial-carcinoma (date last accessed 2 May 2022).
 49. Haugsten EM, Wiedlocha A, Olsnes S, Wesche J. Roles of fibroblast growth factor receptors in carcinogenesis. *Mol Cancer Res*. 2010;8:1439–1452. DOI: 10.1158/1541-7786.MCR-10-0168
 50. Knowles MA, Hurst CD. Molecular biology of bladder cancer: New insights into pathogenesis and clinical diversity. *Nat Rev Cancer*. 2015;15:25–41. DOI: 10.1038/nrc3817
 51. Moss TJ, Qi Y, Xi L, et al. Comprehensive genomic characterization of upper tract urothelial carcinoma. *Eur Urol*. 2017;72:641–649. DOI: 10.1016/j.eururo.2017.05.048
 52. Necchi A, Madison R, Pal SK, et al. Comprehensive genomic profiling of upper-tract and bladder urothelial carcinoma. *Eur Urol Focus*. 2021;7:1339–1346. DOI: 10.1016/j.euf.2020.08.001
 53. Yang K, Yu W, Liu H, et al. Comparison of genomic characterization in upper tract urothelial carcinoma and urothelial carcinoma of the bladder. *Oncologist*. 2021;26:e1395–e1405. DOI: 10.1002/onco.13839
 54. Loriot Y, Necchi A, Park SH, et al. Erdafitinib in locally advanced or metastatic urothelial carcinoma. *N Engl J Med*. 2019;381:338–348. DOI: 10.1056/NEJMoa1817323
 55. Loriot Y, Necchi A, Park SH, et al. Erdafitinib compared with vinorelbine or docetaxel or pembrolizumab in patients (pts) with metastatic or surgically unresectable (M/UR) urothelial carcinoma (UC) and selected fgfr gene alterations (fgfralt): the phase III THOR study. *Ann Oncol*. 2018;29:viu327–viu328. DOI: 10.1093/annonc/mdy283.129
 56. Powles TB, Chistyakov V, Beliakouski V, et al. LBA27 erdafitinib (ERDA) or ERDA plus cetrelimumab (CET) for patients with metastatic or locally advanced urothelial carcinoma (muc) and fibroblast growth factor receptor alterations (fgfra): first phase (ph) II results from the NORSE study. *Ann Oncol*. 2021;32:51303. DOI: 10.1016/j.annonc.2021.08.2103
 57. Tagawa ST, Balar AV, Petrylak DP, et al. TROPHY-U-01: A phase II open-label study of sacituzumab govitecan in patients with metastatic urothelial carcinoma progressing after platinum-based chemotherapy and checkpoint inhibitors. *J Clin Oncol*. 2021;39:2474–2485. DOI: 10.1200/JCO.20.03489
 58. Grivas P, Tagawa ST, Bellmunt J, et al. TROPICS-04: study of sacituzumab govitecan in metastatic or locally advanced unresectable urothelial cancer that has progressed after platinum and checkpoint inhibitor therapy. *J Clin Oncol*. 2021;39(Suppl. 6):TPS498–TPS498. DOI: 10.1200/JCO.2021.39.6_suppl.TPS498
 59. Grivas P, Pouessel D, Park CH, et al. TROPHY-U-01 cohort 3: Sacituzumab govitecan (SG) in combination with pembrolizumab (pembro) in patients (pts) with metastatic urothelial cancer (muc) who progressed after platinum (PLT)-based regimens. *J Clin Oncol*. 2022;40(Suppl. 6):434. DOI: 10.1200/JCO.2022.40.6_suppl.434
 60. Mollica V, Maggio I, Lopez-Beltran A, et al. Combination therapy in advanced urothelial cancer: the role of PARP, HER-2 and mTOR inhibitors. *Expert Rev Anticancer Ther*. 2020;20:755–63. DOI: 10.1080/14737140.2020.1807334
 61. Grivas P, Loriot Y, Morales-Barrera R, et al. Efficacy and safety of rucaparib in previously treated, locally advanced or metastatic urothelial carcinoma from a phase 2, open-label trial (ATLAS). *BMC Cancer*. 2021;21:593. DOI: 10.1186/s12885-021-08085-z
 62. Powles T, Carroll D, Chowdhury S, et al. An adaptive, biomarker-directed platform study of durvalumab in combination with targeted therapies in advanced urothelial cancer. *Nat Med*. 2021;27:793–801. DOI: 10.1038/s41591-021-01317-6
 63. Sadeghi S, Quinn DJ, Dorff T, et al. 6510 phase II trial of pembrolizumab (P) in combination with sephib4-HSA (B4) in previously treated metastatic urothelial carcinoma (muc). *Ann Oncol*. 2021;32:5678. DOI: 10.1016/j.annonc.2021.08.047
 64. Msaouel P, Siefker-Radtke AO, Sweis R, et al. 705MO sitravatinib (sitra) in combination with nivolumab (nivo) demonstrates clinical activity in checkpoint inhibitor (CPI) naïve, platinum-experienced patients (pts) with advanced or metastatic urothelial carcinoma (UC). *Ann Oncol*. 2020;31(Suppl. 4):S556. DOI: 10.1016/j.annonc.2020.08.777
 65. Drakaki A, Rezazadeh Kalebasty A, Lee J-L, et al. Phase Ib/II umbrella trial to evaluate the safety and efficacy of multiple 2L cancer immunotherapy (CI) combinations in advanced/metastatic urothelial carcinoma (muc); MORPHEUS-muc. *J Clin Oncol*. 2020;38(Suppl. 6):TPS591–TPS591. DOI: 10.1200/JCO.2020.38.6_suppl.TPS591
 66. Takai Y, Ikeda W, Ogita H, Rikitake Y. The immunoglobulin-like cell adhesion molecule nectin and its associated protein afadin. *Annu Rev Cell Dev Biol*. 2008;24:309–342. DOI: 10.1146/annurev.cellbio.24.110707.175339
 67. Takai Y, Miyoshi Y, Ikeda W, Ogita H. Nectins and nectin-like molecules: Roles in contact inhibition of cell movement and proliferation. *Nat Rev Mol Cell Biol*. 2008;9:603–615. DOI: 10.1038/nrm2457
 68. Rikitake Y, Takai Y. Interactions of the cell adhesion molecule nectin with transmembrane and peripheral membrane proteins for pleiotropic functions. *Cell Mol Life Sci*. 2008;65:253–263. DOI: 10.1007/s00118-007-7290-9
 69. Chalhita-Eld PM, Satpayev D, Yang P, et al. Enfortumab vedotin antibody-drug conjugate targeting nectin-4 is a highly potent therapeutic agent in multiple preclinical cancer models. *Cancer Res*. 2016;76:3003–3013. DOI: 10.1158/0008-5472.CAN-15-1313
 70. Tomiyama E, Fujita K, Rodriguez Pena MDC, et al. Expression of nectin-4 and PD-L1 in upper tract urothelial carcinoma. *Int J Mol Sci*. 2020;21:5390. DOI: 10.3390/ijms21155390
 71. Hoffman-Censits JH, Lombardo KA, Parimi V, et al. Expression of Nectin-4 in bladder urothelial carcinoma, in morphologic variants, and nonurothelial histotypes. *Appl Immunohistochem Mol Morphol*. 2021;29:619–625. DOI: 10.1097/PAI.0000000000000938
 72. Rosenberg J, Sridhar SS, Zhang J, et al. EV-101: A phase I study of single-agent enfortumab vedotin in patients with nectin-4-positive solid tumors, including metastatic urothelial carcinoma. *J Clin Oncol*. 2020;38:1041–1049. DOI: 10.1200/JCO.19.02044
 73. US Food and Drug Administration. FDA grants accelerated approval to enfortumab vedotin-efjv for metastatic urothelial cancer. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-enfortumab-vedotin-efjv-metastatic-urothelial-cancer (date last accessed 2 May 2022).
 74. Lacouture ME, Patel AB, Rosenberg JE, O’Donnell PH. Management of dermatologic events associated with the nectin-4-directed antibody-drug conjugate enfortumab vedotin. *Oncologist*. 2022;27:e223–e232. DOI: 10.1093/oncoloyagac001
 75. US Food and Drug Administration. FDA grants regular approval to enfortumab vedotin-efjv for locally advanced or metastatic urothelial cancer. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-regular-approval-enfortumab-vedotin-efjv-locally-advanced-or-metastatic-urothelial-cancer (date last accessed 2 May 2022).
 76. Rosenberg JE, Powles T, Sonpavde GP, et al. Long-term outcomes in EV-301: 24-month findings from the phase 3 trial of enfortumab vedotin versus chemotherapy in patients with previously treated advanced urothelial carcinoma. *J Clin Oncol*. 2022;40(Suppl. 16):4516–4516. DOI: 10.1200/JCO.2022.40.16_suppl.4516
 77. Friedlander TW, Milowsky MI, Bilen MA, et al. Study EV-103: Update on durability results and long term outcome of enfortumab vedotin + pembrolizumab in first line locally advanced or metastatic urothelial carcinoma (la/muc). *J Clin Oncol*. 2021;39(Suppl. 15):4528. DOI: 10.1200/JCO.2021.39.15_suppl.4528
 78. Petrylak DP, Flaig TW, Mar N, et al. Study EV-103 cohort H: Antitumor activity of neoadjuvant treatment with enfortumab vedotin monotherapy in patients with muscle-invasive bladder cancer who are cisplatin-ineligible. *J Clin Oncol*. 2022;40(Suppl. 16):4582. DOI: 10.1200/JCO.2022.40.16_suppl.4582
 79. Rosenberg JE, Milowsky M, Ramamurthy C, et al. Study EV-103 cohort K: Antitumor activity of enfortumab vedotin (EV) monotherapy or in combination with pembrolizumab (P) in previously untreated cisplatin-ineligible patients (pts) with locally advanced or metastatic urothelial cancer (la/muc).

- Ann Oncol.* 2022;33(Suppl. 7):S808–69. DOI: 10.1016/annonc/annonc1089
80. McCaffrey JA, Hilton S, Mazumdar M, et al. Phase II trial of docetaxel in patients with advanced or metastatic transitional-cell carcinoma. *J Clin Oncol.* 1997;15:1853–1857. 10.1200/JCO.1997.15.5.1853
 81. Sridhar SS, Blais N, Tran B, et al. Efficacy and safety of nab-paclitaxel vs paclitaxel on survival in patients with platinum-refractory metastatic urothelial cancer: The Canadian Cancer Trials Group BL.12 Randomized Clinical Trial. *JAMA Oncol.* 2020;6:1751–1758. DOI: 10.1001/jamaoncol.2020.3927
 82. Gerullis H, Wawroschek F, Köhne CH, Ecke TH. Vinflunine in the treatment of advanced urothelial cancer: Clinical evidence and experience. *Ther Adv Urol.* 2017;9:28–35. DOI: 10.1177/1756287216677903
 83. US Food and Drug Administration. FDA grants accelerated approval to sacituzumab govitecan for advanced urothelial cancer. www.fda.gov/drugs/resources-information-approved-drugs/fda-grants-accelerated-approval-sacituzumab-govitecan-advanced-urothelial-cancer (date last accessed 4 May 2022).
 84. Makrakis D, Talukder R, Lin GI, et al. Association between sites of metastasis and outcomes with immune checkpoint inhibitors in advanced urothelial carcinoma between sites of metastasis and outcomes with immune checkpoint inhibitors in advanced urothelial carcinoma. *Clin Genitourin Cancer.* 2022;20:e440–52. DOI: 10.1016/j.clgc.2022.06.001
 85. ClinicalTrials.gov. ClinicalTrials.gov. Sacituzumab Govitecan Plus Enfortumab Vedotin for Metastatic Urothelial Carcinoma Progressing on Platinum-based Chemotherapy and PD1/L1 Inhibitors: the Double Antibody Drug Conjugate (DAD) Phase I Trial. ClinicalTrials.gov Identifier: NCT04724018. <https://clinicaltrials.gov/ct2/show/NCT04724018> (date last accessed 4 May 2022).
 86. Jain RK, Kim Y, Rembisz J, et al. Phase II trial of erdafitinib (E) combined with enfortumab vedotin (EV) following platinum and PD-1/L1 inhibitors for metastatic urothelial carcinoma (muc) with FGFR2/3 genetic alterations (gas). *J Clin Oncol.* 2022;40(Suppl. 6):TPS595–TPS595. DOI: 10.1200/JCO.2022.40.6_suppl.TPS595
 87. Teo MY, Ratna N, Regazzi AM, et al. Enfortumab vedotin in FGFR3-mutated advanced urothelial carcinoma. *J Clin Oncol.* 2021;39(Suppl. 6):458–458. DOI: 10.1200/JCO.2021.39.6_suppl.458
 88. Adib E, El Zarif T, Jain RK, et al. FGFR2/3 genomic alterations and response to enfortumab vedotin in metastatic urothelial carcinoma. *BJUJ Compass.* 2022;3:169–172. DOI: 10.1002/bco2.125
 89. Hoimes CJ, Flaig TW, Srinivas S, et al. Study EV-103 cohort L: Evaluating perioperative enfortumab vedotin monotherapy in cis-ineligible muscle-invasive bladder cancer (MIBC) (trial in progress). *J Clin Oncol.* 2022;40(Suppl. 6):TPS587–TPS587.
 90. Hoimes CJ, Bedke J, Lorient Y, et al. KEYNOTE-B15/EV-304: randomized phase 3 study of perioperative enfortumab vedotin plus pembrolizumab versus chemotherapy in cisplatin-eligible patients with muscle-invasive bladder cancer (MIBC). *J Clin Oncol.* 2021;39(Suppl. 15):TPS4587–TPS4587. DOI: 10.1200/JCO.2021.39.15_suppl.TPS4587
 91. Galsky MD, Necchi A, Shore ND, et al. KEYNOTE-905/EV-303: perioperative pembrolizumab or pembrolizumab plus enfortumab vedotin (EV) and cystectomy compared to cystectomy alone in cisplatin-ineligible patients with muscle-invasive bladder cancer (MIBC). *J Clin Oncol.* 2021;39(Suppl. 6):TPS507–TPS507. DOI: 10.1200/JCO.2021.39.6_suppl.TPS507
 92. Chu CE, Sjöström M, Egusa EA, et al. Heterogeneity in *NECTIN4* expression across molecular subtypes of urothelial cancer mediates sensitivity to enfortumab vedotin. *Clin Cancer Res.* 2021;27:5123–5130. 10.1158/1078-0432.CCR-20-4175
 93. Jindal T, Zhang L, Chou J, et al. Biomarkers predictive of response to enfortumab vedotin (EV) treatment in advanced urothelial cancer (auc). *J Clin Oncol.* 2022;40(Suppl. 6):531–531. DOI: 10.1200/JCO.2022.40.6_suppl.531
 94. Bellmunt J, Choueiri TK, Fougeray R, et al. Prognostic factors in patients with advanced transitional cell carcinoma of the urothelial tract experiencing treatment failure with platinum-containing regimens. *J Clin Oncol.* 2010;28:1850–1855. 10.1200/JCO.2009.25.4599
 95. Makrakis D, Talukder R, Dawsey S, et al. Association of time to second-line (2L) immune-checkpoint inhibitors (ICI) and outcomes with ICIs in patients (PTS) with advanced urothelial carcinoma (auc). *J Clin Oncol.* 2022;40(Suppl. 6):505–505. DOI: 10.1200/JCO.2022.40.6_suppl.505
 96. Cabaud O, Berger L, Crompton E, et al. Overcoming resistance to anti-nectin-4 antibody-drug conjugate. *Mol Cancer Ther.* 2022;21:1227–35. DOI: 10.1158/1535-7163.MCT-22-0013
 97. Meszaros P, Hummel I, Klappe K, et al. The function of the ATP-binding cassette (ABC) transporter ABCB1 is not susceptible to actin disruption. *Biochim Biophys Acta.* 2013;1828:340–51. DOI: 10.1016/j.bbmem.2012.10.007
 98. Powles T, Drakaki A, Teoh J-C, et al. A phase 3, randomized, open-label, multicenter, global study of the efficacy and safety of durvalumab (D) + tremelimumab (T) + enfortumab vedotin (EV) or D + EV for neoadjuvant treatment in cisplatin-ineligible muscle-invasive bladder cancer (MIBC) (VOLGA). *J Clin Oncol.* 2022;40(Suppl. 6):TPS579–TPS579. DOI: 10.1200/JCO.2022.40.6_suppl.TPS579
 99. Liu BA, Olson D, Snead K, et al. Abstract 5581: Enfortumab vedotin, an anti-Nectin-4 ADC demonstrates bystander cell killing and immunogenic cell death anti-tumor activity mechanisms of action in urothelial cancers. *Cancer Res.* 2020;80:5581–5581. 10.1158/1538-7445.AM2020-5581