

Management of NF1-related PNs: Approaches for delivering state-of-the-art care

This activity is funded by an independent medical education grant from Alexion and SpringWorks Therapeutics Inc.

The activity is jointly provided by USF Health and touchIME. touchIME is an EBAC® accredited provider.

It is for healthcare professionals outside of the UK.

Disclaimer

- *Unapproved products or unapproved uses of approved products may be discussed by the faculty; these situations may reflect the approval status in one or more jurisdictions*
- *The presenting faculty have been advised by USF Health and touchIME to ensure that they disclose any such references made to unlabeled or unapproved use*
- *No endorsement by USF Health or touchIME of any unapproved products or unapproved uses is either made or implied by mention of these products or uses in USF Health and touchIME activities*
- *USF Health and touchIME accept no responsibility for errors or omissions*

A conversation between:



Dr. Angela Hirbe

Washington University,
St Louis, MO, USA



Prof. Allan Belzberg

Johns Hopkins Medicine,
Baltimore, MD, USA

When should systemic and targeted therapies be considered for adults with NF1-PN?

Prof. Allan Belzberg

Johns Hopkins Medicine,
Baltimore, MD, USA



Dr. Angela Hirbe




Washington University,
St Louis, MO, USA



Clinical burden of NF1-PN in adults



Dutch national expertise center on NF1, retrospective study, 2004–2025
Adult NF1 cohort (N=1,067, 57% female; NF1-PNs, n=501 [47%])



Key

- Disfigurement
- Pain
- Neurological deficits*



≥1 surgery for symptomatic PNs: 46%[†]

- >1 resection for the same tumor: **12%**
- Peri/post-operative complications: **13%[‡]**
- Growth after resection: **36%**

Surgical management comes with post-operative complications, and radiotherapy is associated with a risk of secondary malignancies; the introduction of MEK inhibitors for managing NF1-PNs is a major step forward

*Neurological deficits in the PN-related symptoms graph comprise motor (12%) and sensory (8%) deficits. [†]Partial/debulking was more common than complete resection.

[‡]Hemorrhage, 4%; neurological deficit, 5%; other, 6%.

NF1-PN, neurofibromatosis type 1-associated PN; PN, plexiform neurofibroma; NF1, neurofibromatosis type 1.

Noordhoek CD, et al. *Neuro-Oncol Pract.* 2026; <https://doi.org/10.1093/nop/npag063>.

MEK inhibitors for NF1-PNs in adults

Selumetinib (25 mg/m ² , orally, BID) ¹			Mirdametinib (2 mg/m ² , orally, BID) ²	
Trial	KOMET (NCT04924608), phase III (N=145) Selumetinib (n=71) vs PBO (n=74)		ReNeu (NCT03962543), phase IIb, N=58 Adults; separate pediatric cohort not shown	
Cycles (C)	28 days (continuous)		28 days, 3 weeks on/1 week off (intermittent)	
Efficacy	ORR (%) at C16* <			

Efficacy data should not be compared directly because of differences in the trial designs.

*ORR (CR+PR). CR, disappearance of the target PN; PR, ≥20% decrease in target PN volume vs baseline (both confirmed by consecutive scans 3–6 months after the first response).

†ORR, ≥20% reduction of target PN volume from baseline on consecutive scans during the 24-cycle treatment phase.

‡The observed ORR was significantly higher (p<0.001) than the predefined minimum clinically relevant response rate of 23% for adults.

AE, adverse effect; BID, twice a day; CR, complete response; MEK, mitogen-activated protein kinase; NF1-PN, neurofibromatosis type 1-associated PN;

ORR, overall response rate; PBO, placebo; PN, plexiform neurofibroma; PR partial response.

1. Chen AP, et al. *Lancet*. 2025;405:2217–30; 2. Moertel CL, et al. *J Clin Oncol*. 2024;43:716–29.


How can adverse effects be monitored and managed to maximize therapeutic benefit?

Dr. Angela Hirbe

Washington University,
St Louis, MO, USA



MEK inhibitor-associated adverse effects



How can pain, function and quality of life guide shared decisions in NF1-PN care?

Prof. Allan Belzberg

Johns Hopkins Medicine,
Baltimore, MD, USA



Care provision within the MDT to optimize HRQoL

Drivers of reduced HRQoL in adults with NF1-PNs¹

Chronic
pain

Functional
decline

Risk of MNPST

Psychosocial
stressors*

Persistent or
worsening HRQoL

Community care²⁻⁴

- General patient needs
- Coordination with the specialist NF1 care team
- Patient monitoring and evaluation where access to a specialist team is limited



PCP



Patient



NF1 MDT

MDT led by an NF1 specialist with a nurse coordinator^{4,5}

- Medical and surgical oncology
- Neuro- and plastic surgery
- Pain management
- Genetic counseling
- Social work

HRQoL challenges evolve across developmental stages, underscoring the need for continuous, age-appropriate supportive care, and management within an MDT¹

*Including employment challenges, sexual self-esteem and social isolation.

HRQoL, health-related quality of life; MDT, multidisciplinary team; MNPST, malignant peripheral nerve sheath tumor; NF1, neurofibromatosis type 1;

NF1-PN, NF1-associated plexiform neurofibroma.

1. Armstrong AE, et al. *Neuro Oncol.* 2026;28:847–63; 2. Stewart DR, et al. *Genet Med.* 2018;20:671–82; 3. Abate D, et al. Presented at: The 2026 NF Conference, Denver Co, USA.

June 26–30, 2026. Poster no. 406; 4. Lipinski LJ, et al. *Orphanet J Rare Dis.* 2026;21:219; 5. Lavani S, Brown RM. *J Multidiscip Healthc.* 2024;17:1803–17.